

## **Product Data Sheet**

## **Purified anti-human GM-CSF**

Catalog # / Size: 502201 / 50 µg

502202 / 500 µg

Clone: BVD2-23B6 **Isotype:** Rat IgG2a, κ

Immunogen: E. coli -expressed, recombinant human GM-CSF


Reactivity: Human

**Preparation:** The antibody was purified by affinity chromatography.

**Formulation:** Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.

Concentration: 0.5 mg/ml

**Storage:** The antibody solution should be stored undiluted at 4°C.



## **Applications:**

Applications: ELISA Capture - Quality tested

IP, WB - Reported in the literature

Recommended Usage: Each lot of this antibody is quality control tested by ELISA assay. For ELISA capture applications, a concentration

range of 2-6 µg/ml is recommended. To obtain a linear standard curve, serial dilutions of GM-CSF recombinant protein ranging from 500 to 4 pg/ml are recommended for each ELISA plate. It is recommended that the reagent be

titrated for optimal performance for each application.

Application Notes: ELISA or ELISPOT Capture 1-6: The purified BVD2-23B6 antibody is useful as the capture antibody in a sandwich

ELISA or ELISPOT assay, when used in conjunction with the biotinylated BVD2-21C11 antibody (Cat. No. 502304) as the detecting antibody. The LEAF™ purified antibody is suggested for ELISPOT capture.

Neutralization¹-⁴: The LEAF™ purified antibody (Endotoxin Additional reported applications (for the relevant

formats) include: immunoprecipitation<sup>3</sup> and Western blotting.

Note: For testing human GM-CSF in serum or plasma, BioLegend's ELISA Max™ Sets (Cat. No. 432001 to 432006)

are specially developed and recommended.

 Bacchetta R, et al. 1990. J. Immunol. 144:902.
 Kita H, et al. 1991. J. Exp. Med. 174:745.
 Abrams J, et al. 1992. Immunol. Rev. 127:5. **Application References:** 

Abrams J, et al. 1994. Eosinophils in Allergy and Inflammation. Marcel Dekker New York. p.133.
 Mahanty S, et al. 1992. J. Immunol. 148:3567.
 Klinman D, et al. 1994. Curr. Prot. Immunol.. John Wiley and Sons New York. Unit 6.19.

Description: Granulocyte/macrophage - colony stimulating factor (GM-CSF) is a hematopoietic factor that is produced by activated

T cells, B cells, mast cells, macrophages, fibroblasts and endothelial cells. In addition to supporting colony formation of granulocyte/macrophage progenitors, GM-CSF is a growth factor for erythroid, megakaryocyte and eosinophil progenitors. The BVD2-23B6 antibody reacts with human granulocyte/macrophage - colony stimulating factor (GM-CSF). The BVD2-23B6 antibody can neutralize the bioactivity of natural or recombinant GM-CSF.

Antigen References: 1. Fitzgerald K, et al. Eds. 2001. The Cytokine FactsBook. Academic Press San Diego.

2. Demetri G, et al. 1991. Blood 78:2791.

3. Fan D, et al. 1991. In vivo 5:571.

4. Negrin R, et al. 1992. Adv. Pharmacol. 23:263.

**Related Products: Product** Clone Application

Biotin anti-human GM-CSF BVD2-21C11 ELISA Detection, ELISPOT

Recombinant Human GM-CSF rh GM-CSF HRP Avidin Avidin

TMB Substrate Reagent Set ELISA Assay Diluent (5X)

Detection, ICFC BA, ELISA ELÍSA, ELISPOT, IHC, WB ELISA

**ELISA** 



