

Recombinant Human Interleukin 11 (IL-11)

Publication Number MAN0003562 Rev. 2.00

Catalan Namalan	DYLCOMAE
Catalog Number:	PHC0115
Quantity:	10 µg
Lot Number:	See product label.
Molecular Weight:	19.1 kDa, 179 amino acid residues.
Purity:	≥98% as determined by SDS-PAGE and HPLC analyses.
Biological Activity:	$ED_{50} \le 2$ ng/mL (corresponding to $\ge 5 \times 10^5$ units/mg). The biological activity was determined by measuring the IL-11 dose dependent stimulation of proliferation of murine T11 cells. The optimal concentration should be determined for each specific application. IL-11 is a lymphoid cell growth factor which promotes the growth and survival of B and T cell subsets.
Formulation:	Lyophilized with no additives. Filtered through a 0.2 micron filter prior to lyophilization.
Endotoxin:	<0.1 ng /μg (1 EU/μg)
Production:	Recombinant human IL–11 is produced in <i>E. coli</i> and purified via sequential chromatography.
Reconstitution Recommendation:	We recommend that the vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute recombinant human IL–11 in sterile, distilled water to a concentration of 0.5–1.0 mg/mL. It is important to note that this protein is slow to dissolve. <i>Do not vortex</i> . This solution can be stored at 2°C to 8°C for up to 1 week. For extended storage, it is recommended to further dilute in a buffer containing a carrier protein, such as 0.1% BSA and store in working aliquots at –20°C to –80°C.
Suggested Working Dilutions:	The optimal concentration should be determined for each specific application.
Storage:	The lyophilized protein is stable at room temperature for 1 month, but should be kept at –20°C for long term storage, preferably desiccated. Working aliquots stored with a carrier protein are stable for at least 3 months at –20°C to –80°C. Avoid repeated freeze/thaw cycles.
Expiration Date:	See product label.
References:	Hiraoka, N., E. Allen, I.J. Apel, M.R. Gyetko, and S.J. Weiss (1998) Matrix metalloproteinases regulate neovascularization by acting as pericellular fibrinolysins. Cell 95(3):365–377.

Explanation of Symbols

The symbols present on the product label are explained below:

Symbol	Description
REF	Catalog Number
RUO	Research Use Only
	Use by
***	Manufacturer
[-]	Without, does not contain
from Light	Protect from light
<u> </u>	Directs the user to consult instructions for use (IFU), accompanying the product.

Symbol	Description
LOT	Batch code
IVD	In vitro diagnostic medical device
1	Temperature limitation
EC REP	European Community authorized representative
[+]	With, contains
<u> </u>	Consult accompanying documents

Limited Use Label License: Research Use Only

The purchase of this product conveys to the purchaser the limited, non-transferable right to use the purchased amount of the product only to perform internal research for the sole benefit of the purchaser. No right to resell this product or any of its components is conveyed expressly, by implication, or by estoppel. This product is for internal research purposes only and is not for use in commercial applications of any kind, including, without limitation, quality control and commercial services such as reporting the results of purchaser's activities for a fee or other form of consideration. For information on obtaining additional rights, please contact outlicensing@lifetech.com or Out Licensing, Life Technologies, 5791 Van Allen Way, Carlsbad, California 92008.

For Research Use Only. Caution: Not for human or animal therapeutic or diagnostic use.

Manufacturing site: 7335 Executive Way | Frederick, MD 21704 | Toll Free in USA 800.955.6288

© 2011 Life Technologies Corporation. All rights reserved. The trademarks mentioned herein are the property of Life Technologies Corporation or their respective owners.

For support visit www.lifetechnologies.com/support or email techsupport@lifetech.com

www.lifetechnologies.com Revision Date 13 May 2011

