

DESCRIPTION


Species Reactivity	Human
Specificity	Detects human gp130 in direct ELISAs and Western blots. In Western blots, approximately 10% cross-reactivity with recombinant mouse gp130 is observed. In direct ELISAs, less than 1% cross-reactivity with recombinant human (rh) IL-1 sRI, rhIL-1 sRII, rhIL-2 sRβ, rhIL-2 sRγ, rhIL-4 sR, and rhIL-6 sR is observed.
Source	Polyclonal Goat IgG
Purification	Antigen Affinity-purified
Immunogen	<i>S. frugiperda</i> insect ovarian cell line Sf 21-derived recombinant human gp130 Leu24-Glu619 (Glu619Asp) Accession # P40189
Endotoxin Level	<0.70 EU per 1 µg of the antibody by the LAL method.
Formulation	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
Western Blot	0.1 µg/mL	Recombinant Human gp130 (Catalog # 228-GP)
Neutralization		Measured by its ability to neutralize Oncostatin M/OSM-induced proliferation in the TF-1 human erythroleukemic cell line. Kitamura, T. <i>et al.</i> (1989) <i>J. Cell Physiol.</i> 140 :323. The Neutralization Dose (ND ₅₀) is typically 2-10 µg/mL in the presence of 0.8 ng/mL Recombinant Human Oncostatin M/OSM.

DATA



PREPARATION AND STORAGE

Reconstitution	Reconstitute at 0.2 mg/mL in sterile PBS.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. <ul style="list-style-type: none"> ● 12 months from date of receipt, -20 to -70 °C as supplied. ● 1 month, 2 to 8 °C under sterile conditions after reconstitution. ● 6 months, -20 to -70 °C under sterile conditions after reconstitution.

BACKGROUND

Gp130, the common signal transducing receptor component shared by the functional receptor complexes of the IL-6 family of cytokines, belongs to the class I cytokine receptor family. Binding of IL-6 (IL-11) to either the membrane-anchored or soluble IL-6 R (IL-11 R) initiates the association of IL-6 R (IL-11 R) with gp130 which then undergoes homo-dimerization and signal transduction. With other IL-6 family cytokines, such as LIF and OSM, signal transduction is triggered by the hetero-dimerization of gp130 and LIF R or OSM R.

Gp130 is expressed in all organs examined. Soluble gp130, which apparently arises either from proteolytic cleavage of the membrane-bound receptor or from alternative splicing, has been detected in human serum. At the present time, the *in vivo* functions of soluble gp130 are not clearly understood. In *in vitro* experiments, natural or recombinant soluble gp130 has been shown to have inhibitory effects on OSM and CNTF activities.

References:

1. Narazaki, M. *et al.* (1993) *Blood* **82**:1120.
2. Taga, T. and T. Kishimoto (1997) *Annu. Rev. Immunol.* **15**:797.