

DESCRIPTION

Species Reactivity	Mouse
Specificity	Detects mouse Fcγ RII/RIII (CD32/CD16) in direct ELISAs and Western blots. In direct ELISAs and Western blots, approximately 5% cross-reactivity with recombinant human Fcγ RII is observed.
Source	Polyclonal Goat IgG
Purification	Antigen Affinity-purified
Immunogen	Mouse myeloma cell line NS0-derived recombinant mouse Fcγ RIIB Thr40-Arg217 Accession # P08101
Formulation	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

	Recommended Concentration	Sample
Western Blot	0.1 µg/mL	Recombinant Mouse Fcγ RIIB/CD32b (Catalog # 1460-CD)
Immunocytochemistry	5-15 µg/mL	Immersion fixed mouse splenocytes

PREPARATION AND STORAGE

Reconstitution	Reconstitute at 0.2 mg/mL in sterile PBS.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	<p>Use a manual defrost freezer and avoid repeated freeze-thaw cycles.</p> <ul style="list-style-type: none"> ● 12 months from date of receipt, -20 to -70 °C as supplied. ● 1 month from date of receipt, 2 to 8 °C, reconstituted. ● 6 months from date of receipt, -20 to -70 °C, reconstituted.

BACKGROUND

Fcγ RII, also named CD32, is a low-affinity receptor for the Fc region of complexed IgG. Three genes for human Fcγ RII (A, B and C) and one for mouse (Fcγ RIIB), encoding type I transmembrane proteins, have been identified. The extracellular domain of mouse Fcγ RII shares approximately 95% amino acid sequence identity with that of mouse Fcγ RIII/CD16.