



Anti-mouse TLR6 Antibody

ORDERING INFORMATION

Catalog Number: AF1533

Lot Number: IWC01

Size: 100 µg

Formulation: 0.2 µm filtered solution in PBS with 5% trehalose

Storage: -20° C

Reconstitution: sterile PBS

Specificity: mouse TLR6

Immunogen: NS0-derived rmTLR6 extracellular domain

Ig Type: mouse TLR6 extracellular domain specific goat IgG

Applications: Direct ELISA
Western blot

Preparation

Produced in goats immunized with purified, NS0-derived, recombinant Toll-like Receptor 6 (rmTLR6). Mouse TLR6 specific IgG was purified by mouse TLR6 affinity chromatography. TLR6 belongs to the IL-1 receptor/Toll-like receptor superfamily. These receptors function as pattern recognition receptors for pathogen associated molecular patterns and are implicated in vertebrate immunity.

Formulation

Lyophilized from a 0.2 µm filtered solution in phosphate-buffered saline (PBS) with 5% trehalose.

Reconstitution

Reconstitute with sterile PBS. If 1 mL of PBS is used, the antibody concentration will be 0.1 mg/mL.

Storage

Lyophilized samples are stable for twelve months from date of receipt when stored at -20° C to -70° C. Upon reconstitution, the antibody can be stored at 2° - 8° C for 1 month without detectable loss of activity. Reconstituted antibody can also be aliquotted and stored frozen at -20° C to -70° C **in a manual defrost freezer** for six months without detectable loss of activity. **Avoid repeated freeze-thaw cycles.**

Specificity

This antibody has been selected for its ability to recognize mouse TLR6 in direct ELISAs and western blots.

Applications

Direct ELISA - This antibody can be used at 0.5 - 1.0 µg/mL with the appropriate secondary reagents to detect mouse TLR6. The detection limit for rmTLR6 is approximately 0.1 ng/well. In this format, this antibody shows approximately 30% cross-reactivity with rmTLR1 and less than 1% cross-reactivity with rmTLR2, rhTLR3 and rhTLR4.

Western blot - This antibody can be used at 0.1 - 0.2 µg/mL with the appropriate secondary reagents to detect mouse TLR6. The detection limit for rmTLR6 is approximately 1 ng/lane under non-reducing and reducing conditions.

Optimal dilutions should be determined by each laboratory for each application.