

## Certificate of Analysis

### pGL4.41[*Luc2P*/HSE/Hygro] Vector:

<b>Part No.</b>	<b>Size</b>
E375A	20µg

**Description:** The pGL4.41[*Luc2P*/HSE/Hygro] Vector<sup>(a-e)</sup> contains four copies of a heat shock response element (HSE) that drives transcription of the luciferase reporter gene *Luc2P* (*Photinus pyralis*). *Luc2P* is a synthetically derived luciferase sequence with humanized codon optimization that is designed for high expression and reduced anomalous transcription. The *Luc2P* gene contains hPEST, a protein destabilization sequence, which allows *Luc2P* protein levels to respond more quickly than those of *Luc2* to induction of transcription. The vector backbone contains an ampicillin resistance gene to allow selection in *E. coli* and a gene for hygromycin resistance to allow selection of stably transfected mammalian cell lines.

**Concentration:** 1µg/µl.

**GenBank® Accession Number:** JQ858520.

**Storage Buffer:** The pGL4.41[*Luc2P*/HSE/Hygro] Vector is supplied in 10mM Tris-HCl (pH 7.4), 1mM EDTA.

**Storage Conditions:** See the product information label for storage temperature recommendations. Avoid multiple freeze-thaw cycles and exposure to frequent temperature changes. See the expiration date on the product information label.

**Usage Note:** Concentration gradients may form in frozen products and should be dispersed upon thawing. Mix well prior to use.

Part# 9PIE375  
Printed 5/12



## Quality Control Assays

**Nuclease Assay:** Following incubation of 1µg of the vector in Restriction Enzyme Buffer at 37°C for 16–24 hours, no evidence of nuclease activity is detected by agarose gel electrophoresis.

**Physical Purity:**  $A_{260}/A_{280} \geq 1.80$ ,  $A_{260}/A_{250} \geq 1.05$ .

**Sequence:** The pGL4.41[*Luc2P*/HSE/Hygro] Vector has been completely sequenced and has 100% identity with the published sequence, available at: [www.promega.com/vectors/](http://www.promega.com/vectors/)



**Promega**

### Promega Corporation

2800 Woods Hollow Road	
Madison, WI 53711-5399	USA
Telephone	608-274-4330
Toll Free	800-356-9526
Fax	608-277-2516
Internet	<a href="http://www.promega.com">www.promega.com</a>

### PRODUCT USE LIMITATIONS, WARRANTY, DISCLAIMER

Promega manufactures products for a number of intended uses. Please refer to the product label for the intended use statements for specific products. Promega products contain chemicals which may be harmful if misused. Due care should be exercised with all Promega products to prevent direct human contact.

Each Promega product is shipped with documentation stating specifications and other technical information. Promega products are warranted to meet or exceed the stated specifications. Promega's sole obligation and the customer's sole remedy is limited to replacement of products free of charge in the event products fail to perform as warranted. Promega makes no other warranty of any kind whatsoever, and SPECIFICALLY DISCLAIMS AND EXCLUDES ALL OTHER WARRANTIES OF ANY KIND OR NATURE WHATSOEVER, DIRECTLY OR INDIRECTLY, EXPRESS OR IMPLIED, INCLUDING, WITHOUT LIMITATION, AS TO THE SUITABILITY, PRODUCTIVITY, DURABILITY, FITNESS FOR A PARTICULAR PURPOSE OR USE, MERCHANTABILITY, CONDITION, OR ANY OTHER MATTER WITH RESPECT TO PROMEGA PRODUCTS. In no event shall Promega be liable for claims for any other damages, whether direct, incidental, foreseeable, consequential, or special (including but not limited to loss of use, revenue or profit), whether based upon warranty, contract, tort (including negligence) or strict liability arising in connection with the sale or the failure of Promega products to perform in accordance with the stated specifications.

© 2012 Promega Corporation. All Rights Reserved. Dual-Glo and GloMax are registered trademarks of Promega Corporation.

FuGENE is a registered trademark of Fugent, LLC. GenBank is a registered trademark of the U.S. Department of Health and Human Services. Opti-MEM is a registered trademark of Life Technologies, Inc.

Products may be covered by pending or issued patents or may have certain limitations. Please visit our Web site for more information.

All specifications are subject to change without prior notice.

Product claims are subject to change. Please contact Promega Technical Services or access the Promega online catalog for the most up-to-date information on Promega products.

Signed by:

J. Stevens, Quality Assurance

<sup>(a)</sup>READ THIS FIRST BEFORE OPENING PRODUCT.

The sale of this product and its use are subject to the terms of a limited use label license, the full text of which is available at: [www.promega.com/LULL](http://www.promega.com/LULL). That text must be read by the purchaser prior to opening this product to determine whether the purchaser agrees that all use of the product shall be in accordance with the license terms. If the purchaser is not willing to accept the terms of the limited use label license, Promega is willing to accept the return of the unused product and provide the purchaser with a full refund. However, if the product is opened for any reason, then the purchaser agrees to be bound by the terms of the limited use label license.

<sup>(b)</sup>U.S. Pat. No. 7,728,118.

<sup>(c)</sup>U.S. Pat. No. 5,670,356.


<sup>(d)</sup>U.S. Pat. No. 8,008,006 and European Pat. No. 1341808.

<sup>(e)</sup>The method of recombinant expression of *Coleoptera* luciferase is covered by U.S. Pat. Nos. 5,583,024, 5,674,713 and 5,700,673. A license (from Promega for research reagent products and from The Regents of the University of California for all other fields) is needed for any commercial sale of nucleic acid contained within or derived from this product.

Part# 9PIE375  
Printed in USA. 5/12.

### pGL4.41[*luc2P*/HSE/Hygro] Vector Features List and Map:

HSE response element	285–325
Minimal promoter	371–401
<i>luc2P</i> reporter gene	434–2209
SV40 late poly(A) signal	2249–2470
SV40 early enhancer/promoter	2518–2936
Synthetic hygromycin (Hyg <sup>r</sup> ) coding region	2961–3998
<i>ColE1</i> -derived plasmid replication origin	4394
Synthetic β-lactamase (Amp <sup>r</sup> ) coding region	5185–6045
Synthetic poly(A) signal sequence	4022–4070
Synthetic poly(A) signal/transcriptional pause site	105–258
Reporter Vector primer 3 (RVprimer3) binding region	207–226
Reporter Vector primer 4 (RVprimer4) binding region	4137–4156



Sequence information for the pGL4 Vectors is available online at: [www.promega.com/vectors/](http://www.promega.com/vectors/)

### Example Protocol

In this example protocol, the pGL4.41[*luc2P*/HSE/Hygro] Vector is used to measure activation of the HSE in HepG2 cells upon treatment with 17-AAG or CdCl<sub>2</sub>. The pGL4.75 Vector (encoding *Renilla* luciferase) is used as a normalization control. In designing such experiments, it is important that the chosen cell type can be transfected efficiently and that it expresses the proper components of the signaling pathway of interest in order to generate the biological response. Protocol optimization may be required for your particular cell type and assay conditions.

### Materials to be Supplied by User

- DMEM (Life Technologies Cat.# 11995)
- Complete medium [DMEM supplemented with 10% fetal bovine serum (DMEM/FBS; Life Technologies Cat.# 16000) and 1X NEAA [Life Technologies Cat.# 11140]]
- Dulbecco's PBS (DPBS; Life Technologies Cat.# 14190)
- 0.05% Trypsin-EDTA (Life Technologies Cat.# 25300)
- Charcoal-stripped FBS (Life Technologies Cat.# 126776-011)
- Opti-MEM® I (Life Technologies Cat.# 31985)
- FuGENE® HD Transfection Reagent (Cat.# E2311)
- 17-AAG (17-(Allylamino)-17-demethoxygeldanamycin; Calbiochem Cat.# 100068)
- CdCl<sub>2</sub> (Sigma Cat.# 202908)
- DMSO (Sigma Cat.# D2650)
- Dual-Glo® Luciferase Assay System (Cat.# E2940)
- HepG2 cells
- pGL4.75[*hRLuc*/CMV] Vector (Cat.# E6931)

### Day 1: Plate Cells

1. Grow HepG2 cells in complete medium (DMEM + 10% FBS + 1X NEAA). Wash twice with DPBS and treat with one volume of 0.05% trypsin-EDTA, followed by four volumes of complete medium.
2. Vigorously resuspend the cells by pipetting and allow cell clumps to settle. Remove the cell suspension from any cell clumps, quantify the cells and dilute in complete medium to 1 × 10<sup>5</sup> cells/ml.
3. Plate 100μl per well to a solid, white 96-well plate (Corning Cat.# 3917).
4. Incubate for 24 hours in a 37°C, 5% CO<sub>2</sub> incubator.

### Day 2: Transfection


1. Dilute pGL4.41[*luc2P*/HSE/hygro] and pGL4.75 [*hRLuc*/CMV] *Renilla* luciferase vector constructs in a 10:1 mass ratio, respectively, to 10ng total DNA/μl in Opti-MEM® I.
2. Add FuGENE® HD to a 4.5:1 lipid:DNA ratio. Mix by pipetting. Incubate at room temperature for 20 minutes.
3. Add 10μl transfection complex per well (100ng DNA/well) and incubate for 18 hours in a 37°C, 5% CO<sub>2</sub> incubator.

### Day 3: Medium Replacement and Cell Treatment

1. Resuspend 17-AAG (17-(Allylamino)-17-demethoxygeldanamycin) to 1mM in DMSO. Serially dilute into DMSO to give concentrated stock solutions (1,000X). Serially dilute a 1mM aqueous stock of CdCl<sub>2</sub> into water to give concentrated stock solutions (1,000X). Dilute the 1,000X stocks of 17-AAG and CdCl<sub>2</sub> into DMEM to give 10X stocks.
2. Remove existing medium from cells and replace with 72μl of DMEM + 0.5% charcoal-stripped FBS per well.
3. Add 8μl of the 10X dilutions of 17-AAG or CdCl<sub>2</sub> and incubate for 6 hours in a 37°C, 5% CO<sub>2</sub> incubator.

### Day 4: Luminescence Measurement

1. Remove plates from the 37°C, 5% CO<sub>2</sub> incubator and allow to cool to room temperature for approximately 15 minutes.
2. Add 80μl of the Dual-Glo® Luciferase Assay System detection reagents and measure luminescence following the recommended protocol (Refer to the Dual-Glo® Luciferase Assay System Technical Manual, #TM058 for details).



**Figure 1. Representative data for pGL4.41[*luc2P*/HSE/Hygro] in HepG2 cells upon stimulation with 17-AAG or CdCl<sub>2</sub>.** HepG2 cells were transiently transfected with pGL4.41[*luc2P*/HSE/Hygro] and pGL4.75 and assayed in 96-well format after six hours stimulation with 17-AAG or CdCl<sub>2</sub> as indicated. Firefly luciferase luminescence normalized to the *Renilla* luciferase control is shown. Error bars indicate the S.E.M. for six replicates. Luminescence was detected after addition of Dual-Glo® reagents, using a GloMax® 96 instrument with a 0.5 second integration time. Part# 9PIE375 Printed in USA. 5/12.