

FIP200 (D10D11) Rabbit mAb

✓ 100 µl
(10 western blots)

Orders ■ 877-616-CELL (2355)
orders@cellsignaling.com

Support ■ 877-678-TECH (8324)
info@cellsignaling.com

Web ■ www.cellsignaling.com

New 01/13


For Research Use Only. Not For Use In Diagnostic Procedures.

Applications W, IP Endogenous	Species Cross-Reactivity* H, M, (Mk)	Molecular Wt. 200 kDa	Isotype Rabbit IgG**
-------------------------------------	---	--------------------------	-------------------------


Background: FIP200 (FAK family kinase-interacting protein of 200 kDa) was identified in a two-hybrid screen with the tyrosine kinase Pyk2 and can inhibit Pyk2 kinase activity as well as related family members (1). FIP200 was later independently identified in a multi-drug resistance screen and named RB1CC1 (RB1-inducible coiled-coil 1) due to its induction by cytotoxic stress and RB1 expression regulation (2). FIP200 function has been linked to apoptosis, cell cycle progression, cell growth, and migration (reviewed in 3). FIP200 has also recently been shown to interact with ULK1 and is required for autophagosome formation (4). FIP200 is part of an ULK1 complex along with Atg13 that is regulated by mTOR and is required for starvation induced autophagy (5-7).

Specificity/Sensitivity: FIP200 (D10D11) Rabbit mAb recognizes endogenous levels of total FIP200 protein.


Source/Purification: Monoclonal antibody is produced by immunizing animals with a synthetic peptide corresponding to residues near the carboxy terminus of human FIP200 protein.




Western blot analysis of extracts from various cell lines using FIP200 (D10D11) Rabbit mAb.



Western blot analysis of extracts from A172 cells, untreated (-) or chloroquine-treated (50 µM, overnight; +) using FIP200 (D10D11) Rabbit mAb (upper) or α-Actinin (D6F6) XP® Rabbit mAb #6487 (lower).



Western blot analysis of extracts from 293T cells, mock transfected (-) or transfected with a construct expressing full-length human FIP200 (hFIP200; +), using FIP200 (D10D11) Rabbit mAb.



Immunoprecipitation of FIP200 from JLN-3 cell extracts using Rabbit (DA1E) mAb IgG XP® Isotype Control #3900 (lane 2) or FIP200 (D10D11) Rabbit mAb (lane 3). Lane 1 is 10% input. Western blot analysis was performed using FIP200 (D10D11) Rabbit mAb.

IMPORTANT: For western blots, incubate membrane with diluted antibody in 5% w/v BSA, 1X TBS, 0.1% Tween-20 at 4°C with gentle shaking, overnight.

Applications Key: W—Western IP—Immunoprecipitation IHC—Immunohistochemistry ChIP—Chromatin Immunoprecipitation IF—Immunofluorescence F—Flow cytometry E-P—ELISA-Peptide
Species Cross-Reactivity Key: H—human M—mouse R—rat Hm—hamster Mk—monkey Mi—mink C—chicken Dm—D. melanogaster X—Xenopus Z—zebrafish B—bovine
 Dg—dog Pg—pig Sc—S. cerevisiae Ce—C. elegans Hr—horse All—all species expected Species enclosed in parentheses are predicted to react based on 100% homology.