

Technical Data Sheet

Fixation Buffer

Product Information

Material Number: 554655
Size: 100 mL

Description

BD Cytotfix™ Fixation Buffer is intended to preserve human and rodent lymphoid cells for the subsequent immunofluorescent staining of intracellular cytokines. BD Cytotfix can also be used to preserve the light-scattering characteristics and fluorescence intensities of human and rodent hematopoietic cells that have been stained by immunofluorescence for subsequent flow cytometric analysis.

Preparation and Storage

Store undiluted at 4°C and protected from prolonged exposure to light. Do not freeze.

Application Notes

Recommended Assay Procedure:

BD Cytotfix™ Fixation Buffer can be used to fix unstained cells for subsequent immunofluorescent staining of intracellular cytokines. The suitability of fixing cells for immunofluorescent staining depends on whether the fluorescent antibodies can specifically detect their cognate antigens in a fixed form. With respect to intracellular cytokines, BD Biosciences offers a large panel of conjugated anti-cytokine antibodies that can be successfully used to stain fixed and permeabilized cells. For the staining of antigens expressed on the surface of fixed cells, several fluorescent antibodies directed against mouse cell surface antigens have been identified to be useful.

BD Cytotfix can also be used to fix cells after immunofluorescent staining in order to preserve the light-scattering signals and fluorescent intensities of cells for analysis at a later time. Cell Fixation Buffer may be useful to avoid the capping or shedding of fluorescent antibodies and/or surface antigens during the period before flow cytometric analysis.

Procedure for fixing cells with BD Cytotfix™:

1. Pellet 10^6 suspended cells (e.g., cytokine-producing cells generated by stimulatory culture) by centrifugation (250 - 300 x g) and carefully remove supernatants to avoid cell loss.
2. Add either 200 μ l (for microwell plates) or 500 μ l (for tubes) aliquots of cold DPBS containing protein and NaN₃, gently resuspend cells, pellet, and remove supernatants.
3. Repeat step 2.
4. Add either 100 μ l (for microwell plates) or 250 μ l (for tubes) aliquots of fixation buffer to each cell pellet and resuspend the cells by either pipetting or vortexing. Incubate the cells with fixation buffer for 15 to 30 min at 4°C. (Cell aggregation can be avoided by vortexing prior to the addition of the fixation buffer.)
5. Fixed cells should be washed and suspended in a buffer that contains protein and NaN₃, e.g., either Stain Buffer (FCS) [Cat. No. 554656] or Stain Buffer (BSA) [Cat. No. 554657]. Store the fixed cells at 4°C (protected from light) for subsequent immunofluorescent staining of intracellular cytokines. It is recommended that fixed cell samples be read as soon as possible, i.e., within one week.

For the immunofluorescent staining of intracellular cytokines, cells that have been previously fixed with BD Cytotfix™ can be washed two times in a buffer that contains protein and NaN₃ followed by incubating the cells for at least 10 minutes (4°C) in a buffer containing the cell-permeabilizing agent, saponin. BD Perm/Wash™ buffer (Cat. No. 554723) is ideally suited for this purpose. The fixed and permeabilized cells can then be stained for intracellular cytokines.

Procedure for fixing immunofluorescently-stained cells with BD Cytotfix™:

Cells stained by immunofluorescence for cell surface antigens can be fixed as described above and stored (4°C, protected from light) for subsequent analysis by flow cytometry (or fluorescence microscopy).

Note: BD Cytotfix/Cytoperm™ solution (Cat. No. 554722) and the BD Perm/Wash™ buffer (Cat. No. 554723) are included in BD Cytotfix/Cytoperm Kit (Cat. No. 554714) as well as the BD Cytotfix/Cytoperm Plus Kit with GolgiStop™ (containing monensin; Cat. No. 554715) and BD Cytotfix/Cytoperm Plus Kit with GolgiPlug™ (containing brefeldin A; Cat. No. 555028).

BD Biosciences

bdbiosciences.com

United States	Canada	Europe	Japan	Asia Pacific	Latin America/Caribbean
877.232.8995	866.979.9408	32.2.400.98.95	0120.8555.90	65.6861.0633	55.11.5185.9995

For country contact information, visit bdbiosciences.com/contact

Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton, Dickinson and Company is strictly prohibited.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.

Unless otherwise noted, BD, BD Logo and all other trademarks are property of Becton, Dickinson and Company. © 2015 BD

554655 Rev. 2



Danger: BD Cytotfix™ Fixation Buffer contains 4.21% formaldehyde (w/w).

Hazard statements

Harmful if inhaled.

Causes skin irritation.

Causes serious eye damage.

May cause an allergic skin reaction.

Suspected of causing genetic defects.

May cause cancer. Route of exposure: Inhalative.

May cause respiratory irritation.

Precautionary statements

Wear protective clothing / eye protection.

Wear protective gloves.

Do not breathe mist/vapours/spray.

IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing.

If skin irritation or rash occurs: Get medical advice/attention.

Suggested Companion Products

<u>Catalog Number</u>	<u>Name</u>	<u>Size</u>	<u>Clone</u>
554656	Stain Buffer (FBS)	500 mL	(none)
554657	Stain Buffer (BSA)	500 mL	(none)
554723	Perm/Wash Buffer	100 mL	(none)
554714	BD Cytotfix/Cytoperm™ Fixation/Permeablization Kit	250 Tests	(none)

Product Notices

1. Please refer to www.bdbiosciences.com/pharmingen/protocols for technical protocols.

References

Alaverdi N, Waters JB. *Pharmingen's Hotlines*. 1997:6-15. (Methodology)

BD Biosciences. *Techniques for Immune Function Analysis, Application Handbook 1st Edition*. 2003; Available:

<http://www.bdbiosciences.com/pdfs/manuals/02-8100055-21A1rr.pdf> 2007, Jan. 25. (Methodology)

Lanier LL, Warner NL. Paraformaldehyde fixation of hematopoietic cells for quantitative flow cytometry (FACS) analysis. *J Immunol Methods*. 1981; 47(1):25-30. (Methodology)

Sander B, Andersson J, Andersson U. Assessment of cytokines by immunofluorescence and the paraformaldehyde-saponin procedure. *Immunol Rev*. 1991; 119:65-93. (Methodology)

BD Biosciences

bdbiosciences.com

United States 877.232.8995 Canada 866.979.9408 Europe 32.2.400.98.95 Japan 0120.8555.90 Asia Pacific 65.6861.0633 Latin America/Caribbean 55.11.5185.9995

For country contact information, visit bdbiosciences.com/contact

Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton, Dickinson and Company is strictly prohibited.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.

Unless otherwise noted, BD, BD Logo and all other trademarks are property of Becton, Dickinson and Company. © 2015 BD

554655 Rev. 2

