

Technical Data Sheet



FITC Rat Anti-Mouse Ly-6G

Product Information

Material Number:	561105
Size:	25 µg
Concentration:	0.5 mg/ml
Clone:	1A8
Immunogen:	Ly-6G-transfected EL4J cell line
Isotype:	Rat (LEW) IgG2a, κ
Reactivity:	QC Testing: Mouse
Storage Buffer:	Aqueous buffered solution containing ≤0.09% sodium azide.

Description

The 1A8 antibody reacts with Ly-6G, a 21-25-kDa GPI-anchored protein. In the bone marrow, Ly-6G is expressed on the majority of the largest cells, which are predominantly granulocytes, and not on lymphoid or erythroid cells. In the periphery, it is expressed on granulocytes. The mAb RB6-8C5 (Cat. No. 557445/553123) recognizes both Ly-6G and Ly-6C and blocks the binding of mAb 1A8 to Ly-6G.



Expression of Ly-6G on bone-marrow leukocytes. BALB/c bone-marrow cell suspensions were stained with either FITC-conjugated 1A8 mAb (open histograms) or FITC-conjugated rat IgG2a, κ isotype control mAb R35-95 (Cat. No. 553929, filled histograms), in the presence of Mouse Fc Block™ (purified anti-mouse CD16/CD32, Cat. No. 553141/553142). Non-viable leukocytes were excluded by staining with propidium iodide, and leukocyte subsets were distinguished by their light scatter profiles. Left panel displays lymphoid and erythroid cells, and the right panel displays myeloid cells. Flow cytometry was performed on a FACSCalibur™ (BDIS, San Jose, CA).

Preparation and Storage

The monoclonal antibody was purified from tissue culture supernatant or ascites by affinity chromatography. The antibody was conjugated with FITC under optimum conditions, and unreacted FITC was removed. Store undiluted at 4°C and protected from prolonged exposure to light. Do not freeze.

Application Notes

Application

Flow cytometry	Routinely Tested
----------------	------------------

Recommended Assay Procedure:

For flow cytometry of leukocyte suspensions, we recommend the use of Mouse Fc Block™ (purified anti-mouse CD16/CD32 mAb 2.4G2, Cat. No. 553141/553142).

Suggested Companion Products

Catalog Number	Name	Size	Clone
553141	Purified Rat Anti-Mouse CD16/CD32 (Mouse BD Fc Block™)	0.1 mg	2.4G2
553929	FITC Rat IgG2a, κ Isotype Control	0.25 mg	R35-95

BD Biosciences

bdbiosciences.com

United States	Canada	Europe	Japan	Asia Pacific	Latin America/Caribbean
877.232.8995	888.259.0187	32.53.720.550	0120.8555.90	65.6861.0633	55.11.5185.9995

For country-specific contact information, visit bdbiosciences.com/how_to_order/

Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton Dickinson and Company is strictly prohibited.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.

BD, BD Logo and all other trademarks are the property of Becton, Dickinson and Company. ©2008 BD



Product Notices

1. Since applications vary, each investigator should titrate the reagent to obtain optimal results.
2. Please refer to www.bdbiosciences.com/pharming/protocols for technical protocols.
3. Caution: Sodium azide yields highly toxic hydrazoic acid under acidic conditions. Dilute azide compounds in running water before discarding to avoid accumulation of potentially explosive deposits in plumbing.

References

Fleming TJ, Fleming ML, Malek TR. Selective expression of Ly-6G on myeloid lineage cells in mouse bone marrow. RB6-8C5 mAb to granulocyte-differentiation antigen (Gr-1) detects members of the Ly-6 family. *J Immunol.* 1993; 151(5):2399-2408. (Immunogen)

Fleming TJ, Malek TR. Multiple glycosylphosphatidylinositol-anchored Ly-6 molecules and transmembrane Ly-6E mediate inhibition of IL-2 production. *J Immunol.* 1994; 153(5):1955-1962. (Biology)