

Technical Data Sheet


PE Rat Anti-Mouse CD40

Product Information

Material Number:	561846
Size:	25 µg
Concentration:	0.2 mg/ml
Clone:	3/23
Immunogen:	Mouse CD40 Recombinant Protein
Isotype:	Rat (LOU) IgG2a, κ
Reactivity:	QC Testing: Mouse
Storage Buffer:	Aqueous buffered solution containing ≤0.09% sodium azide.

Description

The 3/23 clone reacts with CD40, a 40-50 kDa glycoprotein expressed on B lymphocytes and other antigen-presenting cells. CD40 has been reported to be transiently expressed on activated CD4+ and CD8+ T cells and in some mouse strains, the 3/23 mAb has been reported to react with 5-10% of T lymphocytes in adult mouse, but not neonatal, spleen. CD40 plays a key role in B-cell growth and differentiation where interactions of CD40 with its ligand, CD154, are involved in the initiation, effector, and memory stages of cell-mediated immune responses. In addition, CD40 has been reported to be involved with the triggering of NK cells and NK-T cells. Ligation of CD40 with the 3/23 antibody has been reported to induce splenic B cells to express the costimulatory molecule CD86 (B7-2). In addition, although the 3/23 antibody by itself is a weak B-cell mitogen, it has been reported to synergize markedly with mitogenic anti-IgM, anti-IgD mAb or IL-4 to promote B-cell proliferation.



Two-color analysis of the expression of CD40 on mouse spleen cells. BALB/c splenocytes were simultaneously stained with FITC Rat anti-Mouse CD4 mAb (Cat. No. 553046) and FITC Rat anti-Mouse CD8a mAb (Cat. No. 553030) and PE Rat anti-Mouse CD40 (right panel). Please note that staining of a T-cell subset by mAb 3/23 has not been consistently observed. Flow cytometry was performed on a BD FACScan™ flow cytometry system.

Preparation and Storage

Store undiluted at 4°C and protected from prolonged exposure to light. Do not freeze.

The monoclonal antibody was purified from tissue culture supernatant or ascites by affinity chromatography.

The antibody was conjugated with R-PE under optimum conditions, and unconjugated antibody and free PE were removed.

Application Notes

Application

Flow cytometry	Routinely Tested
----------------	------------------

Recommended Assay Procedure:

This product has been reported to be compatible with intracellular staining protocols using the BD Cytofix/Cytoperm™ Kit (Cat. No. 554714).

BD Biosciences

bdbiosciences.com

United States	Canada	Europe	Japan	Asia Pacific	Latin America/Caribbean
877.232.8995	888.268.5430	32.53.720.550	0120.8555.90	65.6861.0633	0800.771.7157

For country-specific contact information, visit bdbiosciences.com/how_to_order/

Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton Dickinson and Company is strictly prohibited.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.

BD, BD Logo and all other trademarks are the property of Becton, Dickinson and Company. ©2011 BD



Suggested Companion Products

Catalog Number	Name	Size	Clone
553046	FITC Rat Anti-Mouse CD4	0.1 mg	RM4-5
553030	FITC Rat Anti-Mouse CD8a	0.1 mg	53-6.7
554714	BD Cytotfix/Cytoperm™ Fixation/Permeablization Kit	250 tests	(none)
553930	PE Rat IgG2a, κ Isotype Control	0.1 mg	R35-95
554656	Stain Buffer (FBS)	500 ml	(none)

Product Notices

1. Since applications vary, each investigator should titrate the reagent to obtain optimal results.
2. Please refer to www.bdbiosciences.com/pharming/en/protocols for technical protocols.
3. For fluorochrome spectra and suitable instrument settings, please refer to our Fluorochrome Web Page at www.bdbiosciences.com/colors.
4. Caution: Sodium azide yields highly toxic hydrazoic acid under acidic conditions. Dilute azide compounds in running water before discarding to avoid accumulation of potentially explosive deposits in plumbing.
5. An isotype control should be used at the same concentration as the antibody of interest.

References

Bourgeois C, Rocha B, Tanchot C. A role for CD40 expression on CD8+ T cells in the generation of CD8+ T cell memory. *Science*. 2002; 297(5589):2060-2063. (Biology)

Grewal IS, Flavell RA. CD40 and CD154 in cell-mediated immunity. *Annu Rev Immunol*. 1998; 16:111-135. (Biology)

Hasbold J, Johnson-Leger C, Atkins CJ, Clark EA, Klaus GG. Properties of mouse CD40: cellular distribution of CD40 and B cell activation by monoclonal anti-mouse CD40 antibodies. *Eur J Immunol*. 1994; 24(8):1835-1842. (Immunogen: Activation, (Co)-stimulation)

Hasbold J, Klaus GG. B cells from CBA/N mice do not proliferate following ligation of CD40. *Eur J Immunol*. 1994; 24(1):152-157. (Biology: Activation, (Co)-stimulation)

Klaus GG, Holman M, Hasbold J. Properties of mouse CD40: the role of homotypic adhesion in the activation of B cells via CD40. *Eur J Immunol*. 1994; 24(11):2714-2719. (Biology)

Leifeld L, Trautwein C, Dumoulin FL, Manns MP, Sauerbruch T, Spengler U. Enhanced expression of CD80 (B7-1), CD86 (B7-2), and CD40 and their ligands CD28 and CD154 in fulminant hepatic failure. *Am J Pathol*. 1999; 154(6):1711-1720. (Biology: Immunohistochemistry)

Mahnke K, Becher P, Ricciardi-Castagnoli P, Luger TA, Schawrz T Grabbe S. CD14 is expressed by subsets of murine dendritic cells and upregulated by lipopolysaccharide. In: Ricciardi-Castagnoli P, ed. *Dendritic Cells in Fundamental and Clinical Immunology*. New York: Plenum Press; 1997:145-159. (Biology)

Munder M, Mallo M, Eichmann K, Modolell M. Murine macrophages secrete interferon gamma upon combined stimulation with interleukin (IL)-12 and IL-18: A novel pathway of autocrine macrophage activation. *J Exp Med*. 1998; 187(12):2103-2108. (Biology)

Noelle RJ, Ledbetter JA, Aruffo A. CD40 and its ligand, an essential ligand-receptor pair for thymus-dependent B-cell activation. *Immunol Today*. 1992; 13(11):431-433. (Biology)

Parry SL, Hasbold J, Holman M, Klaus GG. Hypercross-linking surface IgM or IgD receptors on mature B cells induces apoptosis that is reversed by costimulation with IL-4 and anti-CD40. *J Immunol*. 1994; 152(6):2821-2829. (Biology)

Parry SL, Holman MJ, Hasbold J, Klaus GG. Plastic-immobilized anti-mu or anti-delta antibodies induce apoptosis in mature murine B lymphocytes. *Eur J Immunol*. 1994; 24(4):974-979. (Biology)

Tomura M, Yu WG, Ahn HJ, et al. A novel function of Valpha14+CD4+NKT cells: stimulation of IL-12 production by antigen-presenting cells in the innate immune system. *J Immunol*. 1999; 163(1):93-101. (Biology)

Turner JG, Rakhmievich AL, Burdelya L, et al. Anti-CD40 antibody induces antitumor and antimetastatic effects: the role of NK cells. *J Immunol*. 2001; 166(1):89-94. (Biology)