

## SignalSilence® FGF Receptor 4 siRNA II

Cell Signaling  
TECHNOLOGY®✓ 10 µM in 300 µl  
(3 nmol)

**Orders** ■ 877-616-CELL (2355)  
orders@cellsignaling.com

**Support** ■ 877-678-TECH (8324)  
info@cellsignaling.com

**Web** ■ www.cellsignaling.com

New 04/13

For Research Use Only. Not For Use In Diagnostic Procedures.

## Species Cross-Reactivity: H


**Description:** SignalSilence® FGF Receptor 4 siRNA II from Cell Signaling Technology (CST) allows the researcher to specifically inhibit FGF Receptor 4 expression using RNA interference, a method whereby gene expression can be selectively silenced through the delivery of double stranded RNA molecules into the cell. All SignalSilence® siRNA products from CST are rigorously tested in-house and have been shown to reduce target protein expression by western analysis.

**Background:** Fibroblast growth factors (FGFs) produce mitogenic and angiogenic effects in target cells by signaling through cell surface receptor tyrosine kinases. There are four members of the FGF receptor family: FGFR1 (flg), FGFR2 (bek, KGFR), FGFR3, and FGFR4. Each receptor contains an extracellular ligand binding domain, a transmembrane domain, and a cytoplasmic kinase domain (1). Following ligand binding and dimerization, the receptors are phosphorylated at specific tyrosine residues (2). Seven tyrosine residues in the cytoplasmic tail of FGFR1 can be phosphorylated: Tyr463, 583, 585, 653, 654, 730, and 766. Tyr653 and Tyr654 are important for catalytic activity of activated FGFR and are essential for signaling (3). The other phosphorylated tyrosine residues may provide docking sites for downstream signaling components such as Crk and PLCγ (4,5).

**Directions for Use:** CST recommends transfection with 100 nM SignalSilence® FGF Receptor 4 siRNA II 48 to 72 hours prior to cell lysis. For transfection procedure, follow protocol provided by the transfection reagent manufacturer. Please feel free to contact CST with any questions on use.

Each vial contains the equivalent of 100 transfections, which corresponds to a final siRNA concentration of 100 nM per transfection in a 24-well plate with a total volume of 300 µl per well.

**Quality Control:** Oligonucleotide synthesis is monitored base by base through trityl analysis to ensure appropriate coupling efficiency. The oligo is subsequently purified by affinity-solid phase extraction. The annealed RNA duplex is further analyzed by mass spectrometry to verify the exact composition of the duplex. Each lot is compared to the previous lot by mass spectrometry to ensure maximum lot-to-lot consistency.



Western blot analysis of extracts from Huh7 cells, transfected with 100 nM SignalSilence® Control siRNA (Unconjugated) #6568 (-), SignalSilence® FGF Receptor 4 siRNA I #12472 (+), or SignalSilence® FGF Receptor 4 siRNA II, using FGF Receptor 4 (D3B12) XP® Rabbit mAb #8562 (upper) or β-Actin (D6A8) Rabbit mAb #8457 (lower). The FGF Receptor 4 (D3B12) XP® Rabbit mAb confirms silencing of FGF Receptor 4 expression, while the β-Actin (D6A8) Rabbit mAb is used as a loading control.

Entrez-Gene ID #2264  
Swiss-Prot Acc. #P22455

**Storage:** FGF Receptor 4 siRNA II is supplied in RNase-free water. Aliquot and store at -20°C.

Please visit [www.cellsignaling.com](http://www.cellsignaling.com) for a complete listing of recommended companion products.

## Background References:

- (1) Powers, C.J. et al. (2000) *Endocr Relat Cancer* 7, 165-97.
- (2) Reilly, J.F. et al. (2000) *J Biol Chem* 275, 7771-8.
- (3) Mohammadi, M. et al. (1996) *Mol Cell Biol* 16, 977-89.
- (4) Mohammadi, M. et al. (1991) *Mol Cell Biol* 11, 5068-78.
- (5) Larsson, H. et al. (1999) *J Biol Chem* 274, 25726-34.

**Applications Key:** W—Western IP—Immunoprecipitation IHC—Immunohistochemistry ChIP—Chromatin Immunoprecipitation IF—Immunofluorescence F—Flow cytometry E-P—ELISA-Peptide

**Species Cross-Reactivity Key:** H—human M—mouse R—rat Hm—hamster Mk—monkey Mi—mink C—chicken Dm—D. melanogaster X—Xenopus Z—zebrafish B—bovine

Dg—dog Pg—pig Sc—S. cerevisiae Ce—C. elegans Hr—Horse All—all species expected Species enclosed in parentheses are predicted to react based on 100% homology.