Technical Data Sheet

PE Rat Anti-Mouse Ly-6A/E

Product Information

Material Number:	561076
Alternate Name:	Sca-1
Size:	25 µg
Concentration:	0.2 mg/ml
Clone:	E13-161.7
Immunogen:	BALB/c mouse-derived "pre-T" cell hybridoma
Isotype:	Rat (WI) IgG2a, κ
Reactivity:	QC Testing: Mouse
Storage Buffer:	Aqueous buffered solution containing $\leq 0.09\%$ sodium azide.

Description

The E13-161.7 antibody reacts with Ly-6A.2 and Ly-6E.1, which are allelic members of the Ly-6 multigene family. Sca1 (Ly-6A/E), a phosphatidylinositol-anchored protein of about 18 kDa, is expressed on the multipotent hematopoietic stem cell (HSC) in mice with both Ly-6 haplotypes. Sca-1+ HSC are found in the adult bone marrow and fetal liver, but not in the early embryo yolk sac or intraembryonic hematopoietic sites, and can be mobilized to the peripheral blood and spleen in the adult. In mice expressing the Ly-6.2 haplotype (e.g., AKR, C57BL, C57BR, C57L, C58, DBA/2, PL, SJL, SWR, 129), Ly-6A/E is also expressed on distinct subpopulations of bone marrow and peripheral B lymphocytes, myeloid cells, and thymic and peripheral T lymphocytes, on the earliest intrathymic T-cell precursor population, and in several non-hematopoietic tissues. Strains with the Ly-6.1 haplotype (e.g., A, BALB/c, CBA, C3H/He, DBA/1, NZB) have few Ly-6A/E+ resting peripheral lymphocytes, whereas activation of T cells from mice of both Ly-6 haplotypes leads to strong expression of the Sca-1 antigen. Studies with the D7 antibody (Cat. No. 557403) have demonstrated that Ly-6A/E may be involved in the regulation of B and T lymphocyte responses, and it appears to be required for T-cell receptor-mediated T-cell activation. Purified E13-161.7 mAb can block binding of FITC-conjugated D7 antibody (anti-Ly-6A/E, Cat. No. 557405) to mouse splenocytes, but purified mAb D7 (Cat. No. 557403) is unable to block binding of FITC-conjugated E13-161.7 antibody (Cat. No. 553335). Anti-Ly-6A/E (Sca-1) mAb may be used in combination with the Mouse Lineage Panel (Cat. No. 559971) to identify HSC.

Preparation and Storage

The monoclonal antibody was purified from tissue culture supernatant or ascites by affinity chromatography. The antibody was conjugated with R-PE under optimum conditions, and unconjugated antibody and free PE were removed. Store undiluted at 4°C and protected from prolonged exposure to light. Do not freeze.

Application Notes

Application				
Flow cytometry	/tometry Routinely Tested			
Suggested Compar	nion Products			
Catalog Number Name		Size	Clone	
557405	FITC Rat Anti-Mouse Ly-6A/E	0.1 mg	D7	
557403	Purified Rat Anti-Mouse Ly-6A/E	0.1 mg	D7	
553335	FITC Rat Anti-Mouse Ly-6A/E	0.5 mg	E13-161.7	
559971	Biotin Mouse Lineage Panel	1000 tests	(none)	

0.1 mg

R35-95

Product Notices

553930

1. Since applications vary, each investigator should titrate the reagent to obtain optimal results.

PE Rat IgG2a, κ Isotype Control

- 2. Please refer to www.bdbiosciences.com/pharmingen/protocols for technical protocols.
- 3. For fluorochrome spectra and suitable instrument settings, please refer to our Fluorochrome Web Page at www.bdbiosciences.com/colors.
- 4. Caution: Sodium azide yields highly toxic hydrazoic acid under acidic conditions. Dilute azide compounds in running water before discarding to avoid accumulation of potentially explosive deposits in plumbing.

References

Aihara Y, Buhring HJ, Aihara M, Klein J. An attempt to produce "pre-T" cell hybridomas and to identify their antigens. *Eur J Immunol.* 1986; 16(11):1391-1399. (Immunogen)

BD Biosciences

bdbiosciences.com								
United States 877.232.8995	Canada 888.268.5430	Europe 32.53.720.550	Japan 0120.8555.90	Asia Pacific 65.6861.0633	Latin America/Caribbean 0800.771.7157			
For country-spe	cific contact inf	formation, visit	bdbiosciences.co	m/how_to_orde	r/			
Conditions: The in of any patents. BE use of our product product or as a co written authorizat For Research Use (BD, BD Logo and a	formation disclose) Biosciences will m ts. Purchase does n mponent of anoth tion of Becton Dick Only. Not for use in all other trademark	d herein is not to b ot be held responsi ot include or carry er product. Any us inson and Compan diagnostic or there s are the property	e construed as a rec ble for patent infrin any right to resell oi e of this product oth y is strictly prohibite apeutic procedures. of Becton, Dickinsor	ommendation to us gement or other vio r transfer this produ- ner than the permitt ed. Not for resale. n and Company. ©20	e the above product in violation lations that may occur with the ct either as a stand-alone ed use without the express 111 BD			

Kawamoto H, Ohmura K, Katsura Y. Direct evidence for the commitment of hematopoietic stem cells to T, B and myeloid lineages in murine fetal liver. Int Immunol. 1997; 9(7):1011-1019. (Biology)

Marcos MA, Morales-Alcelay S, Godin IE, Dieterlen-Lievre F, Copin SG, Gaspar ML. Antigenic phenotype and gene expression pattern of lymphohemopoietic progenitors during early mouse ontogeny. J Immunol. 1997; 158(6):2627-2637. (Biology)

Osawa M, Nakamura K, Nishi N, et al. In vivo self-renewal of c-Kit+ Sca-1+ Lin(low/-) hemopoietic stem cells. J Immunol. 1996; 156(9):3207-3214. (Biology)