

## Technical Data Sheet



## APC Hamster Anti-Mouse CD11c

## Product Information

<b>Material Number:</b>	561119
<b>Alternate Name:</b>	Cd11c; Itgax; Integrin alpha-X; Integrin $\alpha$ X; Cr4; Complement receptor 4
<b>Size:</b>	25 $\mu$ g
<b>Concentration:</b>	0.2 mg/ml
<b>Clone:</b>	HL3
<b>Immunogen:</b>	C57BL/6 Mouse Intestinal Intraepithelial Lymphocytes
<b>Isotype:</b>	Armenian Hamster IgG1, $\lambda$ 2
<b>Reactivity:</b>	QC Testing: Mouse
<b>Storage Buffer:</b>	Aqueous buffered solution containing protein stabilizer and $\leq 0.09\%$ sodium azide.

## Description

The HL3 monoclonal antibody specifically binds to the integrin  $\alpha$ X chain of gp150, 95 (CD11c/CD18) which is expressed on dendritic cells and CD4<sup>+</sup> CD8<sup>+</sup> intestinal intraepithelial lymphocytes (IEL) and is upregulated on IEL and lymph-node T cells following *in vivo* activation. CD11c is also found on human NK cells. Although its expression on mouse NK cells is not published, we have detected CD11c on mouse splenic NK cells. Cells of the monocyte/macrophage lineage have been reported to express low levels of CD11c. CD11c plays a role in binding of iC3b.



**Expression of CD11c on mouse splenocytes.** C57BL/6 splenocytes were stained with PE-conjugated anti-mouse NK-1.1 mAb PK136 (Cat. No. 557391/553165) and APC-conjugated mAb HL3 (right panel). Note that NK cells show CD11c expression. Flow cytometry was performed on a BD FACSCalibur™ flow cytometry system.

## Preparation and Storage

The monoclonal antibody was purified from tissue culture supernatant or ascites by affinity chromatography.

The antibody was conjugated to APC under optimum conditions, and unconjugated antibody and free APC were removed.

Store undiluted at 4°C and protected from prolonged exposure to light. Do not freeze.

## Application Notes

## Application

Flow cytometry	Routinely Tested
----------------	------------------

## Suggested Companion Products

Catalog Number	Name	Size	Clone
553956	APC Hamster IgG1, $\lambda$ 1 Isotype Control	0.1 mg	G235-2356

## Product Notices

1. Since applications vary, each investigator should titrate the reagent to obtain optimal results.
2. Please refer to [www.bdbiosciences.com/pharmingen/protocols](http://www.bdbiosciences.com/pharmingen/protocols) for technical protocols.
3. This APC-conjugated reagent can be used in any flow cytometer equipped with a dye, HeNe, or red diode laser.

## BD Biosciences

[bdbiosciences.com](http://bdbiosciences.com)

United States	Canada	Europe	Japan	Asia Pacific	Latin America/Caribbean
877.232.8995	888.268.5430	32.53.720.550	0120.8555.90	65.6861.0633	0800.771.7157

For country-specific contact information, visit [bdbiosciences.com/how\\_to\\_order/](http://bdbiosciences.com/how_to_order/)

Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton Dickinson and Company is strictly prohibited.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.

BD, BD Logo and all other trademarks are the property of Becton, Dickinson and Company. ©2011 BD



4. For fluorochrome spectra and suitable instrument settings, please refer to our Fluorochrome Web Page at [www.bdbiosciences.com/colors](http://www.bdbiosciences.com/colors).
5. Caution: Sodium azide yields highly toxic hydrazoic acid under acidic conditions. Dilute azide compounds in running water before discarding to avoid accumulation of potentially explosive deposits in plumbing.
6. Although hamster immunoglobulin isotypes have not been well defined, BD Biosciences Pharmingen has grouped Armenian and Syrian hamster IgG monoclonal antibodies according to their reactivity with a panel of mouse anti-hamster IgG mAbs. A table of the hamster IgG groups, Reactivity of Mouse Anti-Hamster Ig mAbs, may be viewed at [http://www.bdbiosciences.com/pharmingen/hamster\\_chart\\_11x17.pdf](http://www.bdbiosciences.com/pharmingen/hamster_chart_11x17.pdf).

## References

- Barclay NA, Brown MH, Birkeland ML, et al, ed. *The Leukocyte Antigen FactsBook*. San Diego, CA: Academic Press; 1997. (Biology)
- Gao JX, Liu X, Wen J, et al. Differentiation of monocytic cell clones into CD8 alpha+ dendritic cells (DC) suggests that monocytes can be direct precursors for both CD8 alpha+ and CD8 alpha- DC in the mouse. *J Immunol*. 2003; 170(12):5927-5935. (Biology)
- Huleatt JW, Lefrancois L. Antigen-driven induction of CD11c on intestinal intraepithelial lymphocytes and CD8+ T cells in vivo. *J Immunol*. 1995; 154(11):5684-5693. (Immunogen)
- Maraskovsky E, Brasel K, Teepe M, et al. Dramatic increase in the numbers of functionally mature dendritic cells in Flt3 ligand-treated mice: multiple dendritic cell subpopulations identified. *J Exp Med*. 1996; 184(5):1953-1962. (Biology)
- Pulendran B, Lingappa J, Kennedy MK, et al. Developmental pathways of dendritic cells in vivo: distinct function, phenotype, and localization of dendritic cell subsets in FLT3 ligand-treated mice. *J Immunol*. 1997; 159(5):2222-2231. (Biology)