

## Technical Data Sheet


FITC Mouse Anti-Human TCR  $\gamma\delta$ 

## Product Information

Material Number:	561995
Size:	25 $\mu$ g
Concentration:	0.5 mg/ml
Clone:	B1
Isotype:	Mouse IgG1, $\kappa$
Reactivity:	QC Testing: Human
Storage Buffer:	Aqueous buffered solution containing $\leq$ 0.09% sodium azide.

## Description

Recognizes the  $\gamma/\delta$  T-cell receptor (TCR). This receptor complex consists of two disulfide-linked glycoproteins, a  $\gamma$  chain (45-60 kDa) and a  $\delta$  subunit (40-60 kDa).  $\gamma/\delta$  TCR is expressed in less than 10% of human peripheral T cells. The physiological significance of  $\gamma/\delta$ + T cells is still unknown. There is evidence indicating that these cells recognize bacterial ligands and some tumor cells. Reports suggest that  $\gamma/\delta$ + T cells may play a role in the immune reaction during infection and in regulation of pathophysiological autoimmune responses.



Profile of peripheral blood lymphocytes analyzed by flow cytometry.

## Preparation and Storage

Store undiluted at 4°C and protected from prolonged exposure to light. Do not freeze.

The monoclonal antibody was purified from tissue culture supernatant or ascites by affinity chromatography.

The antibody was conjugated with FITC under optimum conditions, and unreacted FITC was removed.

## Application Notes

## Application

Flow cytometry	Routinely Tested
----------------	------------------

## Suggested Companion Products

Catalog Number	Name	Size	Clone
555748	FITC Mouse IgG1, $\kappa$ Isotype Control	100 tests	MOPC-21
555334	PE-Cy™5 Mouse Anti-Human CD3	100 tests	UCHT1

## Product Notices

1. Since applications vary, each investigator should titrate the reagent to obtain optimal results.
2. Please refer to [www.bdbiosciences.com/pharmingen/protocols](http://www.bdbiosciences.com/pharmingen/protocols) for technical protocols.
3. Caution: Sodium azide yields highly toxic hydrazoic acid under acidic conditions. Dilute azide compounds in running water before discarding to avoid accumulation of potentially explosive deposits in plumbing.
4. For fluorochrome spectra and suitable instrument settings, please refer to our Fluorochrome Web Page at [www.bdbiosciences.com/colors](http://www.bdbiosciences.com/colors).

## BD Biosciences

[bdbiosciences.com](http://bdbiosciences.com)

United States	Canada	Europe	Japan	Asia Pacific	Latin America/Caribbean
877.232.8995	888.268.5430	32.53.720.550	0120.8555.90	65.6861.0633	0800.771.7157

For country-specific contact information, visit [bdbiosciences.com/how\\_to\\_order/](http://bdbiosciences.com/how_to_order/)

Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton Dickinson and Company is strictly prohibited.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.

BD, BD Logo and all other trademarks are the property of Becton, Dickinson and Company. ©2011 BD



## References

- Barclay NA, Brown MH, Birkeland ML, et al, ed. *The Leukocyte Antigen FactsBook*. San Diego, CA: Academic Press; 1997. (Biology)
- Breit TM, Wolvers-Tettero IL, van Dongen JJ. Receptor diversity of human T-cell receptor gamma delta expressing cells. *Prog Histochem Cytochem*. 1992; 26(1-4):182-193. (Biology)
- Kabelitz D. Function and specificity of human gamma/delta-positive T cells. *Crit Rev Immunol*. 1992; 11(5):281-303. (Biology)
- Kabelitz D, Pechhold K, Bender A, et al. Activation and activation-driven death of human gamma/delta T cells. *Immunol Rev*. 1991; 120:71-88. (Biology)