

Technical Data Sheet


APC Rat Anti-Mouse IFN- γ

Product Information

Material Number:	562018
Size:	25 μ g
Concentration:	0.2 mg/ml
Clone:	XMG1.2
Immunogen:	Mouse IFN- γ Recombinant Protein
Isotype:	Rat IgG1, κ
Reactivity:	QC Testing: Mouse
Target MW:	15-17 kDa
Storage Buffer:	Aqueous buffered solution containing $\leq 0.09\%$ sodium azide.

Description

The XMG1.2 monoclonal antibody specifically binds to mouse interferon- γ (IFN- γ) protein. IFN- γ is a pleiotropic cytokine, of approximately 15-17 kDa, involved in the regulation of inflammatory and immune responses. It plays an important role in activation, growth, and differentiation of T and B lymphocytes, macrophages, NK cells and other non-hematopoietic cell types. IFN- γ production is associated with the Th1 cell differentiation. The purified form of this antibody has been reported to be a neutralizing antibody.



Expression of IFN- γ by stimulated CD4+ and CD4-C3H spleen cells. Splenocytes from C3H mice were stimulated in culture for 4 hours using PMA (5 ng/ml final concentration; Sigma Cat. #P-8139) and Ionomycin (500 ng/ml final concentration; Sigma Cat. #I0634) in the presence of BD GolgiPlug™ Protein Transport Inhibitor (1 μ l/ml, Cat. No. 555029). The splenocytes were harvested and stained with 0.06 μ g of FITC Rat Anti-Mouse CD4 (Cat. No. 553046), fixed, permeabilized and subsequently stained with 0.12 μ g of APC Rat IgG1 isotype control antibody (Cat. No. 554686, left panel) or with APC Rat Anti-Mouse IFN- γ (middle panel) by using the BD Pharmingen staining protocol. To demonstrate specificity of staining, the binding by the APC-XMG1.2 antibody was blocked by preincubation of the fixed, permeabilized cells with unlabeled XMG1.2 antibody (5.0 μ g; Cat. No. 554409, right panel) prior to staining. The quadrant markers for the bivariate dot plots were set based on the autofluorescence controls and verified using the unlabeled antibody blocking specificity control.

Preparation and Storage

Store undiluted at 4°C and protected from prolonged exposure to light. Do not freeze.

The monoclonal antibody was purified from tissue culture supernatant or ascites by affinity chromatography.

The antibody was conjugated to APC under optimum conditions, and unconjugated antibody and free APC were removed.

Application Notes

Application

Intracellular staining (flow cytometry)	Routinely Tested
---	------------------

Suggested Companion Products

Catalog Number	Name	Size	Clone
554652	MiCK-1 Mouse Cytokine Positive Control Cells	1.0 ml	(none)
554715	BD Cytotfix/Cytoperm Plus Kit (with BD GolgiStop)	250 tests	(none)
554686	APC Rat IgG1, κ Isotype Control	0.1 mg	R3-34
553046	FITC Rat Anti-Mouse CD4	0.1 mg	RM4-5
554409	Purified Rat Anti-Mouse IFN- γ	0.1 mg	XMG1.2
554587	Recombinant Mouse IFN- γ	10 μ g	(none)

BD Biosciences

bdbiosciences.com

United States	Canada	Europe	Japan	Asia Pacific	Latin America/Caribbean
877.232.8995	888.268.5430	32.53.720.550	0120.8555.90	65.6861.0633	0800.771.7157

For country-specific contact information, visit bdbiosciences.com/how_to_order/

Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton Dickinson and Company is strictly prohibited.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.

BD, BD Logo and all other trademarks are the property of Becton, Dickinson and Company. ©2011 BD



Product Notices

1. Since applications vary, each investigator should titrate the reagent to obtain optimal results.
2. Please refer to wwwbdbiosciences.com/pharming/protocols for technical protocols.
3. For fluorochrome spectra and suitable instrument settings, please refer to our Fluorochrome Web Page at wwwbdbiosciences.com/colors.
4. Caution: Sodium azide yields highly toxic hydrazoic acid under acidic conditions. Dilute azide compounds in running water before discarding to avoid accumulation of potentially explosive deposits in plumbing.
5. This APC-conjugated reagent can be used in any flow cytometer equipped with a dye, HeNe, or red diode laser.
6. An isotype control should be used at the same concentration as the antibody of interest.

References

- Abrams JS, Roncarolo MG, Yssel H, Andersson U, Gleich GJ, Silver JE. Strategies of anti-cytokine monoclonal antibody development: immunoassay of IL-10 and IL-5 in clinical samples. *Immunol Rev.* 1992; 127:5-24. (Clone-specific)
- Cherwinski HM, Schumacher JH, Brown KD, Mosmann TR. Two types of mouse helper T cell clone. III. Further differences in lymphokine synthesis between Th1 and Th2 clones revealed by RNA hybridization, functionally monospecific bioassays, and monoclonal antibodies. *J Exp Med.* 1987; 166(5):1229-1244. (Clone-specific)
- Ferrick DA, Schrenzel MD, Mulvania T, Hsieh B, Ferlin WG, Lepper H. Differential production of interferon-gamma and interleukin-4 in response to Th1- and Th2-stimulating pathogens by gamma delta T cells in vivo. *Nature.* 1995; 373(6511):255-257. (Clone-specific: Flow cytometry)
- Hsieh B, Schrenzel MD, Mulvania T, Lepper HD, DiMolfetto-Landon L, Ferrick DA. In vivo cytokine production in murine listeriosis. Evidence for immunoregulation by gamma delta+ T cells. *J Immunol.* 1996; 156(1):232-237. (Clone-specific: Flow cytometry)
- Prussin C, Metcalfe DD. Detection of intracytoplasmic cytokine using flow cytometry and directly conjugated anti-cytokine antibodies. *J Immunol Methods.* 1995; 188(1):117-128. (Methodology: Flow cytometry)
- Sander B, Hoiden I, Andersson U, Moller E, Abrams JS. Similar frequencies and kinetics of cytokine producing cells in murine peripheral blood and spleen. Cytokine detection by immunoassay and intracellular immunostaining. *J Immunol Methods.* 1993; 166(2):201-214. (Clone-specific)
- Vikingsson A, Pederson K, Muller D. Enumeration of IFN-gamma producing lymphocytes by flow cytometry and correlation with quantitative measurement of IFN-gamma. *J Immunol Methods.* 1994; 173(2):219-228. (Clone-specific: Flow cytometry)