

Technical Data Sheet


Purified Rat Anti-Mouse MAdCAM-1

Product Information

Material Number:	550556
Size:	1.0 ml
Concentration:	31.25 µg/ml
Clone:	MECA-367
Immunogen:	Mouse endothelial cells from BALB/c mouse mesenteric and peripheral lymph nodes.
Isotype:	Rat (WI) IgG2a, κ
Reactivity:	QC Testing: Mouse
Storage Buffer:	Aqueous buffered solution containing BSA, goat serum, and ≤0.09% sodium azide.

Description

The MECA-367 antibody reacts with mucosal vascular addressin MAdCAM-1. In the fetus and neonate, MAdCAM-1 is the predominant vascular addressin on the high endothelial venules (HEV) of peripheral lymph nodes. In adult mice, MAdCAM-1 is preferentially expressed in mucosal lymphoid tissues and lamina propria; it is also expressed on sinus-lining cells in the spleen. MAdCAM-1 expression is upregulated on the HEV of peripheral lymph nodes in adult NOD mice⁴ and is involved in the development of diabetes and insulinitis. Furthermore, there is evidence that IFN-γ can induce MAdCAM-1 expression in non-mucosal sites in adult mice. MAdCAM-1 is a predominant ligand for integrin α4β7, a lymphocyte mucosal homing receptor, and a facultative ligand for CD62L (L-selectin). MECA-367 mAb binds to the first domain of MAdCAM-1 and blocks MAdCAM-1-dependent binding *in vitro* and lymphocyte homing to Peyer's patch HEV *in vivo*. Source of the immunogen was endothelial cells from BALB/c mouse mesenteric and peripheral lymph nodes.



Immunohistochemical staining of mouse MadCAM-1:
Frozen sections of normal mouse small intestine were reacted with the anti-MadCAM-1 antibody. Cells of the lamina propria and the mucosal lymphoid tissue expressing MadCAM-1 can be identified by the brown labeling of their cell membranes. Amplification 20X.

Preparation and Storage

Store undiluted at 4°C.

The monoclonal antibody was purified from tissue culture supernatant or ascites by affinity chromatography.

Application Notes

Application

Flow cytometry	Routinely Tested
Immunohistochemistry-frozen	Tested During Development
Immunohistochemistry-formalin (antigen retrieval required)	Not Recommended

Recommended Assay Procedure:

Immunohistochemistry: The MECA-367 antibody specific for mouse MadCAM-1 is recommended to test for immunohistochemical staining of acetone-fixed frozen sections. Tissues tested were mouse spleen, thymus and small intestine. The antibody stains mucosal lymphoid tissue and lamina propria. In the spleen it is expressed in the sinus lining cells. The isotype control recommended for use with this antibody is purified rat IgG2a (Cat. No. 559073). For optimal indirect immunohistochemical staining, the MECA-367 antibody should be titrated (1:10 to 1:50 dilution) and visualized via a three-step staining procedure in combination with polyclonal, biotin conjugated anti-rat Igs (multiple adsorbed) (Cat. No. 559286) as the secondary antibody and Streptavidin-HRP (Cat. No. 550946) together with the DAB detection system (Cat. No. 550880). A detailed protocol of the immunohistochemical procedure is available on our website: www.bdbiosciences.com/support/resources.

BD Biosciences

bdbiosciences.com

United States	Canada	Europe	Japan	Asia Pacific	Latin America/Caribbean
877.232.8995	800.979.9408	32.53.720.550	0120.8555.90	65.6861.0633	55.11.5185.9995

For country contact information, visit bdbiosciences.com/contact

Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton, Dickinson and Company is strictly prohibited.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.

Unless otherwise noted, BD, BD Logo and all other trademarks are property of Becton, Dickinson and Company. © 2011 BD



Suggested Companion Products

Catalog Number	Name	Size	Clone
559073	Purified Rat IgG2a κ Isotype Control	0.25 mg	R35-95
559286	Biotin Goat Anti-Rat Ig	0.5 mg	Polyclonal
550880	DAB Substrate Kit	500 tests	(none)
550946	Streptavidin HRP	50 ml	(none)
559148	Antibody Diluent for IHC	125 ml	(none)

Product Notices

1. Since applications vary, each investigator should titrate the reagent to obtain optimal results.
2. Caution: Sodium azide yields highly toxic hydrazoic acid under acidic conditions. Dilute azide compounds in running water before discarding to avoid accumulation of potentially explosive deposits in plumbing.
3. Source of all serum proteins is from USDA inspected abattoirs located in the United States.
4. This antibody has been developed for the immunohistochemistry application. However, a routine immunohistochemistry test is not performed on every lot. Researchers are encouraged to titrate the reagent for optimal performance.
5. An isotype control should be used at the same concentration as the antibody of interest.
6. Please refer to www.bdbiosciences.com/pharming/en/protocols for technical protocols.

References

Berg EL, McEvoy LM, Berlin C, Bargatze RF, Butcher EC. L-selectin-mediated lymphocyte rolling on MAdCAM-1. *Nature*. 1993; 366(6456):695-698. (Biology)

Berlin C, Berg EL, Briskin MJ, et al. Alpha 4 beta 7 integrin mediates lymphocyte binding to the mucosal vascular addressin MAdCAM-1. *Cell*. 1993; 74(1):185-195. (Biology)

Briskin MJ, McEvoy LM, Butcher EC. MAdCAM-1 has homology to immunoglobulin and mucin-like adhesion receptors and to IgA1. *Nature*. 1993; 363(6428):461-464. (Biology)

Kraal G, Schornagel K, Streeter PR, Holzmann B, Butcher EC. Expression of the mucosal vascular addressin, MAdCAM-1, on sinus-lining cells in the spleen. *Am J Pathol*. 1995; 147(3):763-771. (Biology)

Streeter PR, Berg EL, Rouse BT, Bargatze RF, Butcher EC. A tissue-specific endothelial cell molecule involved in lymphocyte homing. *Nature*. 1988; 331(6151):41-46. (Immunogen)

BD Biosciences

bdbiosciences.com

United States	Canada	Europe	Japan	Asia Pacific	Latin America/Caribbean
877.232.8995	800.979.9408	32.53.720.550	0120.8555.90	65.6861.0633	55.11.5185.9995

For country contact information, visit bdbiosciences.com/contact

Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton, Dickinson and Company is strictly prohibited.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.

Unless otherwise noted, BD, BD Logo and all other trademarks are property of Becton, Dickinson and Company. © 2011 BD

