Technical Data Sheet

PE Mouse Anti-Human LAIR-1

Product Information


550811 **Material Number:** 100 tests Size: 20 µl Vol. per Test: DX26 Clone: Mouse IgG1, κ Isotype: QC Testing: Human Reactivity:

Workshop:

Aqueous buffered solution containing BSA and ≤0.09% sodium azide. Storage Buffer:

Description

eacts with leukocyte associated immunoglobulin-like receptor-1 (LAIR-1), a 32 kDa transmembrane glycoprotein with a single immunoglobulin-like domain and a cytoplasmic tail containing two immune receptor tyrosine-based inhibitory motifs. It is expressed on the majority of human peripheral blood mononuclear leukocytes. LAIR-1 recruits SHP-1 and SHP-2 tyrosine phosphatases upon activation, and cross-linking of the LAIR-1 molecule on NK cells results in strong inhibition of NK cell-mediated cytotoxicity. Although it is structurally related to human killer cell inhibitory receptors, LAIR-1 does not appear to recognize MHC class I molecules and thus represents a novel MHC class I-independent mechanism of NK cell regulation. Other studies report other roles of LAIR-1 in cell differentiation.



Profile of peripheral blood lymphocytes analyzed by flow cytometry

Preparation and Storage

The monoclonal antibody was purified from tissue culture supernatant or ascites by affinity chromatography. The antibody was conjugated with R-PE under optimum conditions, and unconjugated antibody and free PE were removed. Store undiluted at 4° C and protected from prolonged exposure to light. Do not freeze.

Application Notes

Application

 (
Flow cytometry	Routinely Tested	

Suggested Companion Products

Catalog Number	Name	Size	Clone
555749	PE Mouse IgG1, κ Isotype Control	100 tests	MOPC-21

BD Biosciences

www.bdbiosciences.com

United States Canada Europe 32.53.720.550 0120.8555.90 877.232.8995 888.259.0187 65.6861.0633 55.11.5185.9995 For country-specific contact information, visit www.bdbiosciences.com/how_to_order/

Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation Conditions: The information disclosed nerein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton Dickinson and Company is strictly prohibited.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.

BD, BD Logo and all other trademarks are the property of Becton, Dickinson and Company. ©2007 BD



Product Notices

- 1. This reagent has been pre-diluted for use at the recommended Volume per Test. We typically use 1 X 10e6 cells in a 100-μl experimental sample (a test).
- 2. Since applications vary, each investigator should titrate the reagent to obtain optimal results.
- 3. Please refer to www.bdbiosciences.com/pharmingen/protocols for technical protocols.
- 4. Caution: Sodium azide yields highly toxic hydrazoic acid under acidic conditions. Dilute azide compounds in running water before discarding to avoid accumulation of potentially explosive deposits in plumbing.
- 5. Source of all serum proteins is from USDA inspected abattoirs located in the United States.

References

Meyaard L, Adema GJ, Chang C. LAIR-1, a novel inhibitory receptor expressed on human mononuclear leukocytes. 1997; 7(2):283-290.(Biology)
Poggi A, Tomasello E, Ferrero E, Zocchi MR, Moretta L. p40/LAIR-1 regulates the differentiation of peripheral blood precursors to dendritic cells induced by granulocyte-monocyte colony-stimulating factor. *Eur J Immunol.* 1998; 28(7):2086-2091.(Biology)

Xu M, Zhao R, Zhao ZJ. Identification and characterization of leukocyte-associated Ig-like receptor-1 as a major anchor protein of tyrosine phosphatase SHP-1 in hematopoietic cells. *J Biol Chem.* 2000; 275(23):17440-17446.(Biology)

550811 Rev. 4 Page 2 of 2