

Technical Data Sheet

APC Goat Anti-Mouse Ig (Multiple Adsorption)**Product Information**

Material Number:	550826
Size:	0.1 mg
Concentration:	0.2 mg/ml
Clone:	Polyclonal
Isotype:	Goat Ig
Reactivity:	QC Testing: Mouse
Storage Buffer:	Aqueous buffered solution containing ≤0.09% sodium azide.

Preparation and Storage

The polyclonal antibody was purified from antiserum by negative adsorption and affinity chromatography.

The antibody was conjugated to APC under optimum conditions, and unconjugated antibody and free APC were removed.

Store undiluted at 4°C and protected from prolonged exposure to light. Do not freeze.

The antibody solution was passed through solid-phase immunoabsorbent gels to minimize cross-reactivity with rat, human, bovine and horse serum proteins.

Application Notes**Application**

Flow cytometry	Routinely Tested
----------------	------------------

Recommended Assay Procedure:

As a second-step reagent for staining rat spleen and bone marrow leukocytes, it is reactive with mouse primary antibodies having the IgG1, IgG2a, IgG2b, IgG3, IgM, and IgA isotypes. Minimal background staining of rat leukocytes occurs in the absence of primary antibody. In addition, it stains mouse peripheral B lymphocytes with little non-specific staining of other splenic leukocytes. Therefore, it is useful as a primary reagent in immunofluorescent staining of mouse B cells and antibody-producing cells and as a secondary reagent for staining rat leukocytes with mouse Ig primary antibodies. However, we have observed that the reactivity of polyclonal second-step antibodies to mouse or rat IgM may be reduced after adsorption against Ig of rat or mouse, respectively. Because this anti-mouse Ig antibody was adsorbed with rat Ig, it may be weakly reactive with some mouse IgM primary antibodies. In those cases, we recommend biotinylated anti-mouse IgM mAb R6-60.2 (Cat. No. 553406) or anti-mouse Ig κ light chain mAb 187.1 (Cat. No. 559750), followed by Streptavidin-APC (Cat. No. 554067).

Suggested Companion Products

Catalog Number	Name	Size	Clone
553406	Biotin Rat Anti-Mouse IgM	0.5 mg	R6-60.2
559750	Biotin Rat Anti-Mouse Ig, κ Light Chain	0.5 mg	187.1
554067	APC Streptavidin	0.1 mg	(none)

Product Notices

1. Since applications vary, each investigator should titrate the reagent to obtain optimal results.
2. Please refer to www.bdbiosciences.com/pharmingen/protocols for technical protocols.
3. For fluorochrome spectra and suitable instrument settings, please refer to our Fluorochrome Web Page at www.bdbiosciences.com/colors.
4. Caution: Sodium azide yields highly toxic hydrazoic acid under acidic conditions. Dilute azide compounds in running water before discarding to avoid accumulation of potentially explosive deposits in plumbing.

BD Biosciences

bdbiosciences.com

United States 877.232.8995 Canada 888.268.5430 Europe 32.53.720.550 Japan 0120.8555.90 Asia Pacific 65.6861.0633 Latin America/Caribbean 0800.771.7157

For country-specific contact information, visit bdbiosciences.com/how_to_order/

Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton Dickinson and Company is strictly prohibited.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.

BD, BD Logo and all other trademarks are the property of Becton, Dickinson and Company. ©2011 BD

