Technical Data Sheet

Purified Mouse Anti-Human CD8

Product Information

Material Number: 551346 Size: 0.1 mg **Concentration:** 0.5 mg/ml Clone: G42-8


Isotype: Mouse IgG2a, κ QC Testing: Human Reactivity:

Workshop:

Storage Buffer: Aqueous buffered solution containing ≤0.09% sodium azide.

Description

Reacts with the a subunit (32 kDa) of the two-chain complex. CD8 molecule binds to HLA class I molecules during interaction of CD8+T cells with antigen- presenting cells or with target cells. CD8 is expressed on T cytotoxic/ suppressor cell populations. G42-8 stains approximately 13-48% of peripheral blood lymphocytes and 80% of thymocytes, as well as a subset of NK cells. This antibody is useful in detecting CD8+ T cells when using DimerX I reagents (Cat. No. 551263).



Profile of peripheral blood lymphocytes analyzed by flow cytometry. Second step staining with Cat. No. 555988

Preparation and Storage

The monoclonal antibody was purified from tissue culture supernatant or ascites by affinity chromatography. Store undiluted at 4° C.

Application Notes

Application

Elever existence et mis	Doutingly Tostad
I Flow cytometry	Routinely Tested

Suggested Companion Products

Catalog Number	Name	Size	Clone	
555571	Purified Mouse IgG2a, κ Isotype Control	0.1 mg	G155-178	
551263	DimerX I: Recombinant Soluble Dimeric Human HLA-A2:Ig Fusion Protein	0.05 mg	(none)	
555988	FITC Goat Anti-Mouse IgG/IgM	0.5 mg	Polyclonal	

BD Biosciences

www.bdbiosciences.com

Europe 877.232.8995 888.259.0187 32.53.720.550 0120.8555.90 65.6861.0633 55.11. For country-specific contact information, visit www.bdbiosciences.com/how_to_order/ 55.11.5185.9995

Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation Conditions: The information disclosed nerein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton Dickinson and Company is strictly prohibited.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.

BD, BD Logo and all other trademarks are the property of Becton, Dickinson and Company. @2007 BD



Product Notices

- 1. Since applications vary, each investigator should titrate the reagent to obtain optimal results.
- 2. Please refer to www.bdbiosciences.com/pharmingen/protocols for technical protocols.
- 3. Caution: Sodium azide yields highly toxic hydrazoic acid under acidic conditions. Dilute azide compounds in running water before discarding to avoid accumulation of potentially explosive deposits in plumbing.

References

Schlossman S, Boumell L, et al, ed. Leucocyte Typing V. New York: Oxford University Press; 1995.(Biology)

551346 Rev. 3 Page 2 of 2