# Technical Data Sheet Purified Mouse Anti-Pig IgG

## **Product Information**

Material Number:	552554
Size:	0.5 mg
Concentration:	0.5 mg/ml
Clone:	F007-1241
Immunogen:	Purified pig IgG
Isotype:	Mouse (BALB/c) IgG1, ĸ
Reactivity:	QC Testing: Pig
Storage Buffer:	Aqueous buffered solution containing $\leq 0.09\%$ sodium azide.

### Description

The F007-1241 antibody reacts with pig IgG, but not with pig IgM. The reactivity with individual pig IgG subclasses (IgG1, IgG2a, IgG2b, IgG3, IgG4) was not determined. It does not cross-react with mouse, rat, hamster, or rabbit immunoglobulins. The F007-1241 antibody does have some cross-reactivity with dog immunoglobulins.

### Preparation and Storage

The monoclonal antibody was purified from tissue culture supernatant or ascites by affinity chromatography.

# Store undiluted at 4° C.

Application	Notes
-------------	-------

Application				
	ELISA Capture	Routinely Tested		

### **Recommended Assay Procedure:**

We recommend using the purified F007-1241 mAb for capture in sandwich pig IgG ELISA in combination with biotinylated anti-pig IgG F007-1241 mAb (Cat. No. 552555) for detection.

# **Suggested Companion Products**

Catalog Number	Name	Size	Clone
552555	Biotin Mouse Anti-Pig IgG	0.2 mg	F007-1241

### **Product Notices**

- 1. Since applications vary, each investigator should titrate the reagent to obtain optimal results.
- 2. Please refer to www.bdbiosciences.com/pharmingen/protocols for technical protocols.
- 3. Caution: Sodium azide yields highly toxic hydrazoic acid under acidic conditions. Dilute azide compounds in running water before discarding to avoid accumulation of potentially explosive deposits in plumbing.
- 4. Sodium azide is a reversible inhibitor of oxidative metabolism; therefore, antibody preparations containing this preservative agent must not be used in cell cultures nor injected into animals. Sodium azide may be removed by washing stained cells or plate-bound antibody or dialyzing soluble antibody in sodium azide-free buffer. Since endotoxin may also affect the results of functional studies, we recommend the NA/LE (No Azide/Low Endotoxin) antibody format, if available, for in vitro and in vivo use.

BD Biosciences							
www.bdbioscie	ences.com						
United States	Canada	Europe	Japan	Asia Pacific	Latin America/Caribbean		
877.232.8995	888.259.0187	32.53.720.550	0120.8555.90	65.6861.0633	55.11.5185.9995		
For country-spe	cific contact inf	formation, visit <b>v</b>	www.bdbioscien	ces.com/how_to	_order/		
Conditions: The in: of any patents. BD use of our product product or as a cor written authorizat For Research Use C	formation disclose Biosciences will n s. Purchase does n mponent of anoth ion of Becton Dick Only. Not for use in	d herein is not to b ot be held responsil ot include or carry a er product. Any use inson and Compan diagnostic or there	e construed as a rec ble for patent infrin any right to resell o e of this product oth y is strictly prohibite apeutic procedures.	ommendation to use gement or other vio r transfer this produc er than the permitte d. Not for resale.	e the above product in violation lations that may occur with the ct either as a stand-alone ed use without the express		

BD, BD Logo and all other trademarks are the property of Becton, Dickinson and Company. ©2007 BD