

Technical Data Sheet


Biotin Mouse Anti-Mouse Vβ 8.1, 8.2 TCR

Product Information

Material Number:	553184
Size:	0.25 mg
Concentration:	0.5 mg/ml
Clone:	MR5-2
Immunogen:	C57BL/6 mouse helper T-cell clone OI6
Isotype:	Mouse (C57L) IgG2a, κ
Reactivity:	QC Testing: Mouse
Storage Buffer:	Aqueous buffered solution containing ≤0.09% sodium azide.

Description

The MR5-2 antibody reacts with the Vβ 8.1 and Vβ 8.2 T-cell Receptors (TCR), but not the Vβ 8.3 TCR, of mice having the *b* haplotype (e.g., A, AKR, BALB/c, CBA/Ca, CBA/J, C3H/He, C57BL, C58, DBA/1, DBA/2) of the *Tcrb* gene complex. The *Tcrb-Vβ* subfamily gene loci are deleted in mice having the *a* (e.g., C57BR, C57L, SJL, SWR) or *c* (e.g., RIII) haplotype. Vβ 8.1 TCR-bearing T lymphocytes are clonally eliminated in mice expressing superantigen encoded by the *Mtv-7* (*Mls-1a*, *Mlsa*), provirus (e.g., AKR, CBA/J, C58, DBA/2), and activation or elimination of Vβ 8.1 TCR-expressing T cells by this determinant is partially dependent upon presentation by I-E. *Mtv-43* (e.g., MA/MyJ), *Mtv-44* (e.g., NZW), and/or exogenous MMTV-SW superantigens also cause incomplete elimination of Vβ 8.1 TCR-bearing T cells. In addition to expression on conventional T lymphocytes, Vβ 8.2 is the predominant β chain of the TCR on NK-T cells. Staphylococcal enterotoxin B, in association with antigen presenting cells expressing I-A and/or I-E, stimulates lymphocytes bearing Vβ 8 TCR and selectively eliminates those T cells *in vivo*. Plate-bound MR5-2 antibody activates Vβ 8.1 or 8.2 TCR-bearing T lymphocytes.



Two-color analysis of the expression of Vβ 8.1, 8.2 TCR on peripheral T lymphocytes. C57BL/6 lymph node cells were incubated simultaneously with biotinylated MR5-2, PE-conjugated RM4-5 (anti-CD4, Cat. No. 553048/553049), and PE-conjugated 53-6.7 (anti-CD8a, Cat. No. 553032/553033) monoclonal antibodies, followed by Avidin-FITC (Cat. No. 554057). Flow cytometry was performed on a BD FACScan™ Flow Cytometry System.

Preparation and Storage

The monoclonal antibody was purified from tissue culture supernatant or ascites by affinity chromatography. The antibody was conjugated with biotin under optimum conditions, and unreacted biotin was removed. Store undiluted at 4°C and protected from prolonged exposure to light. Do not freeze.

Application Notes

Application

Flow cytometry	Routinely Tested
Immunohistochemistry-frozen	Tested During Development

Recommended Assay Procedure:

For flow cytometry of cell suspensions from peripheral lymphoid tissues, it is recommended that multicolor staining be performed to distinguish T lymphocytes from non-T cells.

BD Biosciences

[bdbiosciences.com](http://bdbiosciences.com)

United States	Canada	Europe	Japan	Asia Pacific	Latin America/Caribbean
877.232.8995	888.259.0187	32.53.720.550	0120.8555.90	65.6861.0633	55.11.5185.9995

For country-specific contact information, visit [bdbiosciences.com/how\\_to\\_order/](http://bdbiosciences.com/how_to_order/)

Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton Dickinson and Company is strictly prohibited.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.

BD, BD Logo and all other trademarks are the property of Becton, Dickinson and Company. ©2008 BD



## Suggested Companion Products

Catalog Number	Name	Size	Clone
553048	PE Rat Anti-Mouse CD4	0.1 mg	RM4-5
553032	PE Rat Anti-Mouse CD8a	0.1 mg	53-6.7
554057	Avidin FITC	0.5 mg	(none)
553455	Biotin Mouse IgG2a, $\kappa$ Isotype Control	0.25 mg	G155-178

## Product Notices

1. Since applications vary, each investigator should titrate the reagent to obtain optimal results.
2. Please refer to [www.bdbiosciences.com/pharming/en/protocols](http://www.bdbiosciences.com/pharming/en/protocols) for technical protocols.
3. Caution: Sodium azide yields highly toxic hydrazoic acid under acidic conditions. Dilute azide compounds in running water before discarding to avoid accumulation of potentially explosive deposits in plumbing.

## References

Behlke MA, Chou HS, Huppi K, Loh DY. Murine T-cell receptor mutants with deletions of beta-chain variable region genes. *Proc Natl Acad Sci U S A*. 1986; 83(3):767-771.(Biology)

Bendelac A. Mouse NK1+ T cells. *Curr Opin Immunol*. 1995; 7(3):367-374.(Biology)

Fairchild S, Rosenwasser OA, Dyson PJ, Tomonari K. Tcrb-V3+ T-cell deletion and a new mouse mammary tumor provirus, Mtv-44. *Immunogenetics*. 1992; 36(3):189-194.(Biology)

Haqqi TM, Banerjee S, Anderson GD, David CS. RIII S/J (H-2r). An inbred mouse strain with a massive deletion of T cell receptor V beta genes. *J Exp Med*. 1989; 169(6):1903-1909.(Biology)

Hodes RJ, Abe R. Mouse endogenous superantigens: Mls and Mls-like determinants encoded by mouse retroviruses. In: Coligan JE, Kruisbeek AM, Margulies DH, Shevach EM, Strober W, ed. *Current Protocols in Immunology*. New York: John Wiley & Sons; 1996:A.1F.1-A.1F.5.(Biology)

Hugo P, Kappler JW, Godfrey DI, Marrack PC. Thymic epithelial cell lines that mediate positive selection can also induce thymocyte clonal deletion. *J Immunol*. 1994; 152(3):1022-1031.(Biology)

Kanagawa O. Antibody-mediated activation of T cell clones as a method for screening hybridomas producing antibodies to the T cell receptor. *J Immunol Methods*. 1988; 110(2):169-178.(Immunogen)

Kruisbeek AM, Shevach EM. Proliferative assays for T cell function. In: Coligan J, Kruisbeek AM, Margulies D, Shevach EM, Strober W, ed. *Current Protocols in Immunology*. New York: John Wiley and Sons; 1991:3.12.1-3.12.14.(Biology)

Shinohara K, Ikarashi Y, Maruoka H, et al. Functional and phenotypical characteristics of hepatic NK-like T cells in NK1.1-positive and -negative mouse strains. *Eur J Immunol*. 1999; 29(6):1871-1878.(Biology)

Tomonari K, Fairchild S. Positive and negative selection of Tcrb-V6+ T cells. *Immunogenetics*. 1992; 36(4):230-237.(Biology)

White J, Herman A, Pullen AM, Kubo R, Kappler JW, Marrack P. The V beta-specific superantigen staphylococcal enterotoxin B: stimulation of mature T cells and clonal deletion in neonatal mice. *Cell*. 1989; 56(1):27-35.(Biology)