

Technical Data Sheet


PE Rat Anti-Mouse Vα 11.1, 11.2[b,d] TCR

Product Information

Material Number:	553223
Size:	0.1 mg
Concentration:	0.2 mg/ml
Clone:	RR8-1
Immunogen:	Mouse T-Cell Clone B10
Isotype:	Rat IgG2b, κ
Reactivity:	QC Testing: Mouse
Storage Buffer:	Aqueous buffered solution containing ≤0.09% sodium azide.

Description

The RR8-1 antibody reacts with the Vα 11.1 and Vα 11.2, but not Vα 11.3 T-cell Receptors (TCR) of mice having the *b* (e.g., C57BL) and *d* (e.g., DBA/1, DBA/2, NZW) haplotypes of *Tcra* gene complex. RR8-1 antibody does not react with strains having the *a* (e.g., A, AKR, BALB/c, CBA, C3H/He) or *c* (e.g., NZB, SJL, SWR, NOD) *Tcra* haplotypes. Plate-bound RR8-1 antibody activates Vα 11.1, 11.2[b,d] TCR-bearing T lymphocytes.



Two-color analysis of the expression of Vα 11.1, 11.2[b,d] TCR on peripheral lymphocytes. C57BL/6 lymph node cells were incubated simultaneously with PE-conjugated RR8-1, FITC-conjugated RM4-5 (anti-CD4, Cat. No. 553046/553047), and FITC-conjugated 53-6.7 (anti-CD8a, Cat. No. 553030/553031) monoclonal antibodies. Flow cytometry was performed on a FACScan™ (BDIS, San Jose, CA).

Preparation and Storage

The monoclonal antibody was purified from tissue culture supernatant or ascites by affinity chromatography. The antibody was conjugated with R-PE under optimum conditions, and unconjugated antibody and free PE were removed. Store undiluted at 4°C and protected from prolonged exposure to light. Do not freeze.

Application Notes

Application

Flow cytometry	Routinely Tested
----------------	------------------

Recommended Assay Procedure:

For flow cytometry of cell suspensions from peripheral lymphoid tissues, it is recommended that multicolor staining be performed to distinguish T lymphocytes from non-T cells. This antibody conjugate has been tested by immunofluorescent staining (≤ 1 µg/million cells) with flow cytometric analysis.

Suggested Companion Products

Catalog Number	Name	Size	Clone
553046	FITC Rat Anti-Mouse CD4	0.1 mg	RM4-5
553030	FITC Rat Anti-Mouse CD8a	0.1 mg	53-6.7
553989	PE Rat IgG2b, κ Isotype Control	0.1 mg	A95-1

Product Notices

- Since applications vary, each investigator should titrate the reagent to obtain optimal results.

BD Biosciences

bdbiosciences.com

United States	Canada	Europe	Japan	Asia Pacific	Latin America/Caribbean
877.232.8995	888.268.5430	32.53.720.550	0120.8555.90	65.6861.0633	0800.771.7157

For country-specific contact information, visit bdbiosciences.com/how_to_order/

Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton Dickinson and Company is strictly prohibited.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.

BD, BD Logo and all other trademarks are the property of Becton, Dickinson and Company. ©2011 BD



2. Caution: Sodium azide yields highly toxic hydrazoic acid under acidic conditions. Dilute azide compounds in running water before discarding to avoid accumulation of potentially explosive deposits in plumbing.
3. For fluorochrome spectra and suitable instrument settings, please refer to our Fluorochrome Web Page at www.bdbiosciences.com/colors.
4. Please refer to www.bdbiosciences.com/pharmingen/protocols for technical protocols.

References

Jameson SC, Nakajima PB, Brooks JL, Heath W, Kanagawa O, Gascoigne NR. The T cell receptor V alpha 11 gene family. Analysis of allelic sequence polymorphism and demonstration of J alpha region-dependent recognition by allele-specific antibodies. *J Immunol.* 1991; 147(9):3185-3193. (Immunogen)