

Technical Data Sheet

Purified Rat Anti-Mouse IgD

Product Information

Material Number:	553438
Size:	0.5 mg
Concentration:	0.5 mg/ml
Clone:	11-26c.2a
Immunogen:	Not reported
Isotype:	Rat IgG2a, κ
Reactivity:	QC Testing: Mouse
Storage Buffer:	Aqueous buffered solution containing $\leq 0.09\%$ sodium azide.

Description

The 11-26c.2a antibody reacts with mouse immunoglobulin D of all *Igh-C* haplotypes (e.g., IgDa, IgDb, IgDe), and it does not react with other immunoglobulin isotypes. Although 11-26c.2a mAb binds membrane IgD expressed on the splenic B-cell surface with high affinity, it does not induce proliferation of spleen B cells *in vitro*. *In vivo* injection of 11-26c.2a antibody does not have any effect on activation of mature B cells, as determined by Ia expression.

Preparation and Storage

The monoclonal antibody was purified from tissue culture supernatant or ascites by affinity chromatography.

Store undiluted at 4° C.

Application Notes

Application

Flow cytometry	Routinely Tested
Immunohistochemistry-frozen	Reported

Suggested Companion Products

Catalog Number	Name	Size	Clone
553927	Purified Rat IgG2a, κ Isotype Control	0.5 mg	R35-95
554016	FITC Goat Anti-Rat Ig	0.5 mg	Polyclonal

Product Notices

1. Since applications vary, each investigator should titrate the reagent to obtain optimal results.
2. Please refer to www.bdbiosciences.com/pharmingen/protocols for technical protocols.
3. Caution: Sodium azide yields highly toxic hydrazoic acid under acidic conditions. Dilute azide compounds in running water before discarding to avoid accumulation of potentially explosive deposits in plumbing.

References

Hamilton AM, Lehuen A, Kearney JF. Immunofluorescence analysis of B-1 cell ontogeny in the mouse. *Int Immunol*. 1994; 6(3):355-361.(Biology)
 Ishihara K, Wood WJ Jr, Wall R, et al. Multiple B29 containing complexes on murine B lymphocytes. Common and stage-restricted Ig-associated polypeptide chains. *J Immunol*. 1993; 150(6):2253-2262.(Biology)
 Nitschke L, Kosco MH, Kohler G, Lamers MC. Immunoglobulin D-deficient mice can mount normal immune responses to thymus-independent and -dependent antigens. *Proc Natl Acad Sci U S A*. 1993; 90(5):1887-1891.(Biology)

BD Biosciences

www.bdbiosciences.com

United States 877.232.8995 Canada 888.259.0187 Europe 32.53.720.550 Japan 0120.8555.90 Asia Pacific 65.6861.0633 Latin America/Caribbean 55.11.5185.9995

For country-specific contact information, visit www.bdbiosciences.com/how_to_order/

Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton Dickinson and Company is strictly prohibited.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.

BD, BD Logo and all other trademarks are the property of Becton, Dickinson and Company. ©2007 BD

