

Technical Data Sheet

Biotin Mouse Anti-Mouse IgM[b]

Product Information

Material Number:	553519
Alternate Name:	Igh-6b
Size:	0.5 mg
Concentration:	0.5 mg/ml
Clone:	AF6-78
Immunogen:	C57BL/10 mouse splenocytes
Isotype:	Mouse (BALB/c) IgG1, κ
Reactivity:	QC Testing: Mouse
Storage Buffer:	Aqueous buffered solution containing $\leq 0.09\%$ sodium azide.

Description

The AF6-78 antibody reacts specifically with mouse IgM of *Igh-C[b]* and related haplotypes (e.g., C57BL/6, C57BL/10, SJL, AKR, NZB). It does not react with IgM of *Igh-C[a]* or related haplotypes (e.g., BALB/c, C58, CBA, C3H/Bi, C3H/He, DBA/1, DBA/2, PL, RIII). Cross-reaction with IgM of *Igh-C[e]* haplotype (e.g., A/J) has been observed. AF6-78 antibody does not react with free μ heavy chain in vitro or in the cytoplasm of pre-B lymphocytes, which lack Ig light chain. It has not been shown to stimulate B-cell proliferation.

This antibody is routinely tested by ELISA and flow cytometric analysis. Other applications were tested at BD Biosciences Pharmingen during antibody development only or reported in the literature.

Preparation and Storage

The monoclonal antibody was purified from tissue culture supernatant or ascites by affinity chromatography.

The antibody was conjugated with biotin under optimum conditions, and unreacted biotin was removed.

Store undiluted at 4° C and protected from prolonged exposure to light. Do not freeze.

Application Notes

Application

ELISA	Routinely Tested
Flow cytometry	Routinely Tested

Recommended Assay Procedure:

Biotin-conjugated AF6-78 mAb may be used as a primary or secondary reagent in immunofluorescent staining.

Suggested Companion Products

Catalog Number	Name	Size	Clone
550615	Biotin Mouse IgG1 κ Isotype Control	0.25 mg	MOPC-31C
554057	Avidin FITC	0.5 mg	(none)

Product Notices

1. Since applications vary, each investigator should titrate the reagent to obtain optimal results.
2. Please refer to www.bdbiosciences.com/pharmingen/protocols for technical protocols.
3. Caution: Sodium azide yields highly toxic hydrazoic acid under acidic conditions. Dilute azide compounds in running water before discarding to avoid accumulation of potentially explosive deposits in plumbing.

References

- Stall AM. Mouse immunoglobulin allotypes. In: Herzenberg LA, Weir DM, Blackwell C, ed. *Weir's Handbook of Experimental Immunology*. Blackwell Science Publishers; 1996:27.1-27.16.(Clone-specific)
- Stall AM, Loken MR. Allotypic specificities of murine IgD and IgM recognized by monoclonal antibodies. *J Immunol*. 1984; 132(2):787-795.(Immunogen)

BD Biosciences

www.bdbiosciences.com

United States 877.232.8995 Canada 888.259.0187 Europe 32.53.720.550 Japan 0120.8555.90 Asia Pacific 65.6861.0633 Latin America/Caribbean 55.11.5185.9995

For country-specific contact information, visit www.bdbiosciences.com/how_to_order/

Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton Dickinson and Company is strictly prohibited.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.

BD, BD Logo and all other trademarks are the property of Becton, Dickinson and Company. ©2007 BD

