

Technical Data Sheet

FITC Mouse Anti-Mouse I-A[b]

Product Information

Material Number:	553551
Size:	0.5 mg
Concentration:	0.5 mg/ml
Clone:	AF6-120.1
Immunogen:	Mouse C57BL/10J
Isotype:	Mouse (BALB/c) IgG2a, κ
Reactivity:	QC Testing: Mouse
Storage Buffer:	Aqueous buffered solution containing $\leq 0.09\%$ sodium azide.

Description

The AF6-120.1 antibody reacts with the I-A[b] MHC class II alloantigen. It cross-reacts with cells from mice of the H-2[k] and H-2[u] haplotypes. Reactivity with other haplotypes (e.g., *d, f, g7, p, q, r, s*) has not been observed.

This antibody is routinely tested by flow cytometric analysis. Other applications were tested at BD Biosciences Pharmingen during antibody development only or reported in the literature.

Preparation and Storage

The monoclonal antibody was purified from tissue culture supernatant or ascites by affinity chromatography.

The antibody was conjugated with FITC under optimum conditions, and unreacted FITC was removed.

Store undiluted at 4° C and protected from prolonged exposure to light. Do not freeze.

Application Notes

Application

Flow cytometry	Routinely Tested
----------------	------------------

Suggested Companion Products

Catalog Number	Name	Size	Clone
553456	FITC Mouse IgG2a, κ Isotype Control	0.25 mg	G155-178

Product Notices

1. Since applications vary, each investigator should titrate the reagent to obtain optimal results.
2. Please refer to www.bdbiosciences.com/pharmingen/protocols for technical protocols.
3. Caution: Sodium azide yields highly toxic hydrazoic acid under acidic conditions. Dilute azide compounds in running water before discarding to avoid accumulation of potentially explosive deposits in plumbing.

References

Beck BN, Buerstedde JM, Krco CJ, Nilson AE, Chase CG, McKean DJ. Characterization of cell lines expressing mutant I-Ab and I-Ak molecules allows the definition of distinct serologic epitopes on A alpha and A beta polypeptides. *J Immunol.* 1986; 136(8):2953-2961.(Clone-specific: Flow cytometry)

Cohn LE, Glimcher LH, Waldmann RA, et al. Identification of functional regions on the I-Ab molecule by site-directed mutagenesis. *Proc Natl Acad Sci U S A.* 1986; 83(3):747-751.(Clone-specific: Flow cytometry)

Hattori M, Buse JB, Jackson RA, et al. The NOD mouse: recessive diabetogenic gene in the major histocompatibility complex. *Science.* 1986; 231(4739):733-735. (Biology)

Nabozny GH, Baisch JM, Cheng S, et al. HLA-DQ8 transgenic mice are highly susceptible to collagen-induced arthritis: a novel model for human polyarthritis. *J Exp Med.* 1996; 183(1):27-37.(Clone-specific: Flow cytometry)

Wall KA, Lorber MI, Loken MR, McClatchey S, Fitch FW. Inhibition of proliferation of Mls- and Ia-reactive cloned T cells by a monoclonal antibody against a determinant shared by I-A and I-E. *J Immunol.* 1983; 131(3):1056-1064.(Clone-specific: Flow cytometry)

BD Biosciences

bdbiosciences.com

United States	Canada	Europe	Japan	Asia Pacific	Latin America/Caribbean
877.232.8995	888.259.0187	32.53.720.550	0120.8555.90	65.6861.0633	55.11.5185.9995

For country-specific contact information, visit bdbiosciences.com/how_to_order/

Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton Dickinson and Company is strictly prohibited.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.

BD, BD Logo and all other trademarks are the property of Becton, Dickinson and Company. ©2006 BD



BD

BD Biosciences