

## Technical Data Sheet

## PE Mouse Anti-Mouse H-2D[d]

## Product Information

<b>Material Number:</b>	553580
<b>Size:</b>	0.1 mg
<b>Concentration:</b>	0.2 mg/ml
<b>Clone:</b>	34-2-12
<b>Immunogen:</b>	(C57BL/6 x DBA/2)F1 mouse splenocytes
<b>Isotype:</b>	Mouse (C3H) IgG2a, $\kappa$
<b>Reactivity:</b>	QC Testing: Mouse
<b>Storage Buffer:</b>	Aqueous buffered solution containing $\leq 0.09\%$ sodium azide.

## Description

The 34-2-12 antibody (also known as 34-2-12S) recognizes the  $\alpha 3$  domain of the H-2D[d]. The binding of the antibody to its epitope is independent of the  $\alpha 1$  and  $\alpha 2$  domains and  $\beta 2$  microglobulin. It cross-reacts with cells of the C3H.LG/Ckc strain. Reactivity with other haplotypes (eg, *b, f, k, p, q, r, s*) has not been observed. Soluble mAb 34-2-12 blocks binding of the Ly-49A-expressing T lymphoma EL4 to immobilized H-2D[d]. However, further studies utilizing this mAb indicate that the  $\alpha 3$  domain is not involved in the interaction between Ly-49A, or Ly-49G2, and H-2D[d].

This antibody is routinely tested by flow cytometric analysis. Other applications were tested at BD Biosciences Pharmingen during antibody development only or reported in the literature.

## Preparation and Storage

The monoclonal antibody was purified from tissue culture supernatant or ascites by affinity chromatography.

The antibody was conjugated with R-PE under optimum conditions, and unconjugated antibody and free PE were removed by gel filtration chromatography.

Store undiluted at 4° C and protected from prolonged exposure to light. Do not freeze.

## Application Notes

## Application

Flow cytometry	Routinely Tested
----------------	------------------

## Suggested Companion Products

Catalog Number	Name	Size	Clone
553457	PE Mouse IgG2a, $\kappa$ Isotype Control	0.1 mg	G155-178

## Product Notices

1. Since applications vary, each investigator should titrate the reagent to obtain optimal results.
2. Please refer to [www.bdbiosciences.com/pharmingen/protocols](http://www.bdbiosciences.com/pharmingen/protocols) for technical protocols.
3. Caution: Sodium azide yields highly toxic hydrazoic acid under acidic conditions. Dilute azide compounds in running water before discarding to avoid accumulation of potentially explosive deposits in plumbing.

## References

- Daniels BF, Karlhofer FM, Seaman WE, Yokoyama WM. A natural killer cell receptor specific for a major histocompatibility complex class I molecule. *J Exp Med.* 1994; 180(2):687-692.(Clone-specific: Blocking)
- Evans GA, Margulies DH, Shykind B, Seidman JG, Ozato K. Exon shuffling: mapping polymorphic determinants on hybrid mouse transplantation antigens. *Nature.* 1982; 300(5894):755-757.(Biology)
- Kane KP. Ly-49 mediates EL4 lymphoma adhesion to isolated class I major histocompatibility complex molecules. *J Exp Med.* 1994; 179(3):1011-1015.(Clone-specific: Blocking)
- Karlhofer FM, Ribaldo RK, Yokoyama WM. MHC class I alloantigen specificity of Ly-49+ IL-2-activated natural killer cells. *Nature.* 1992; 358(6381):66-70.(Biology)
- Mason LH, Ortaldo JR, Young HA, Kumar V, Bennett M, Anderson SK. Cloning and functional characteristics of murine large granular lymphocyte-1: a member of the Ly-49 gene family (Ly-49G2). *J Exp Med.* 1995; 182(2):293-303.(Biology)
- McCluskey J, Bluestone JA, Coligan JE, Maloy WL, Margulies DH. Serologic and T cell recognition of truncated transplantation antigens encoded by in vitro deleted class I major histocompatibility genes. *J Immunol.* 1986; 136(4):1472-1481.(Clone-specific: Immunoprecipitation)

## BD Biosciences

[www.bdbiosciences.com](http://www.bdbiosciences.com)

United States	Canada	Europe	Japan	Asia Pacific	Latin America/Caribbean
877.232.8995	888.259.0187	32.53.720.550	0120.8555.90	65.6861.0633	55.11.5185.9995

For country-specific contact information, visit [www.bdbiosciences.com/how\\_to\\_order/](http://www.bdbiosciences.com/how_to_order/)

Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton Dickinson and Company is strictly prohibited.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.

BD, BD Logo and all other trademarks are the property of Becton, Dickinson and Company. ©2007 BD



McCluskey J, Germain RN, Margulies DH. Cell surface expression of an in vitro recombinant class II/class I major histocompatibility complex gene product. *J Immunol.* 1985; 40(2):247-257.(Clone-specific: Immunoprecipitation)

Otten GR, Bikoff E, Ribaldo RK, Kozlowski S, Margulies DH, Germain RN. Peptide and beta 2-microglobulin regulation of cell surface MHC class I conformation and expression. *J Immunol.* 1992; 148(12):3723-3732.(Biology)

Ozato K, Mayer NM, Sachs DH. Monoclonal antibodies to mouse major histocompatibility complex antigens. *Transplantation.* 1982; 34(3):113-120.(Immunogen: Cytotoxicity)