Technical Data Sheet Biotin Mouse Anti-Rat IgG1

Product Information	
Material Number:	553890
Size:	0.5 mg
Concentration:	0.5 mg/ml
Clone:	RG11/39.4
Immunogen:	Pooled rat IgG1
Isotype:	Mouse (SJL) IgG2a, ĸ
Reactivity:	QC Testing: Rat
Storage Buffer:	Aqueous buffered solution containing ${\leq}0.09\%$ sodium azide.

Description

The RG11/39.4 antibody reacts specifically with the Fc region of rat IgG1. It does not react with other Ig isotypes.

This antibody is routinely tested by ELISA and flow cytometric analysis. Other applications were tested at BD Biosciences Pharmingen during antibody development only or reported in the literature.

Preparation and Storage

The monoclonal antibody was purified from tissue culture supernatant or ascites by affinity chromatography. The antibody was conjugated with biotin under optimum conditions, and unreacted biotin was removed. Store undiluted at 4° C and protected from prolonged exposure to light. Do not freeze.

Application Notes

A	Application					
	ELISA	Routinely Tested				
	Flow cytometry	Routinely Tested				

Recommended Assay Procedure:

For the sandwich rat IgG1 ELISA, biotinylated mAb RG11/39.4 is optimal for detection with purified anti-rat IgG1 mAb B46-2 (available by special order) for capture. Biotin-conjugated RG11/39.4 mAb may be used as a secondary reagent in immunofluorescent staining RG11/39.4 antibody is effective for detection of cell-surface or intracellular Ig by immunofluorescent staining with flow cytometric analysis. For flow cytometric detection of intracytoplasmic IgG1, we recommend FITC-conjugated RG11/39.4 mAb (Cat. No. 553892). For immunohistochemical staining, we recommend the use of biotinylated anti-rat IgG1/2a mAb G28-5 in our special formulation for immunohistochemistry, Cat. No. 550325.

Suggested Companion Products

Catalog Number Name		Size	Clone
553892	FITC Mouse Anti-Rat IgG1	0.5 mg	RG11/39.4

Product Notices

- 1. Since applications vary, each investigator should titrate the reagent to obtain optimal results.
- 2. Please refer to www.bdbiosciences.com/pharmingen/protocols for technical protocols.
- 3. Caution: Sodium azide yields highly toxic hydrazoic acid under acidic conditions. Dilute azide compounds in running water before discarding to avoid accumulation of potentially explosive deposits in plumbing.

References

Springer TA, Bhattacharya A, Cardoza JT, Sanchez-Madrid F. Monoclonal antibodies specific for rat IgG1, IgG2a, and IgG2b subclasses, and kappa chain monotypic and allotypic determinants: reagents for use with rat monoclonal antibodies. *Hybridoma*. 1982; 1(3):257-273.(Immunogen: ELISA)

BD Biosciences

www.bdbiosciences.com								
United States	Canada	Europe	Japan	Asia Pacific	Latin America/Caribbean			
877.232.8995	888.259.0187	32.53.720.550	0120.8555.90	65.6861.0633	55.11.5185.9995			
For country-specific contact information, visit www.bdbiosciences.com/how_to_order/								
Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton Dickinson and Company is strictly prohibited. For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale. BD, BD Logo and all other trademarks are the property of Becton, Dickinson and Company. ©2007 BD								