Technical Data Sheet

Biotin Mouse Anti-Rat IgG2a

Product Information

 Material Number:
 553894

 Size:
 0.5 mg

 Concentration:
 0.5 mg/ml

 Clone:
 RG7/1.30

 Immunogen:
 Rat Pooled IgG

 Isotype:
 Mouse (SJL) IgG2b, κ

 Reactivity:
 QC Testing: Rat

Storage Buffer: Aqueous buffered solution containing ≤0.09% sodium azide.

Description

The Rg7/1.30 antibody reacts specifically with the Fc region of rat IgG2a. It does not react with other Ig isotypes.

This antibody is routinely tested by flow cytometric analysis and ELISA. Other applications were tested at BD Biosciences Pharmingen during antibody development only or reported in the literature.

Preparation and Storage

The monoclonal antibody was purified from tissue culture supernatant or ascites by affinity chromatography. The antibody was conjugated with biotin under optimum conditions, and unreacted biotin was removed. Store undiluted at 4° C and protected from prolonged exposure to light. Do not freeze.

Application Notes

Application

ipprettion .		
Flow cytometry	Routinely Tested	
ELISA Detection	Routinely Tested	

Recommended Assay Procedure:

For the sandwich rat IgG2a ELISA, biotinylated mAb RG7/1.30 is optimal for detection with purified anti-rat IgG2a mAb B46-7 (Cat. No. 553918) for capture. Biotin-conjugated RG7/1.30 mAb may be used as a secondary reagent in immunofluorescent staining. RG7/1.30 antibody is effective for detection of cell-surface or intracellular Ig by immunofluorescent staining with flow cytometric analysis. For flow cytometric detection of intracytoplasmic Ig2a, we recommend FITC-conjugated RG7/1.30 mAb (Cat. No. 553896). For immunohistochemical staining, we recommend the use of biotinylated anti-rat IgG1/2a mAb G28-5 in our special formulation for immunohistochemistry, Cat. No. 550325.

Suggested Companion Products

Catalog Number	Name	Size	Clone	
553896	FITC Mouse Anti-Rat IgG2a	0.5 mg	RG7/1.30	
553918	Purified Mouse Anti-Rat IgG2a	0.5 mg	B46-7	
550325	Biotin Mouse Anti-Rat IgG1/2a	1.0 ml	G28-5	

Product Notices

- 1. Since applications vary, each investigator should titrate the reagent to obtain optimal results.
- 2. Please refer to www.bdbiosciences.com/pharmingen/protocols for technical protocols.
- Caution: Sodium azide yields highly toxic hydrazoic acid under acidic conditions. Dilute azide compounds in running water before discarding to avoid accumulation of potentially explosive deposits in plumbing.

References

Springer TA, Bhattacharya A, Cardoza JT, Sanchez-Madrid F. Monoclonal antibodies specific for rat IgG1, IgG2a, and IgG2b subclasses, and kappa chain monotypic and allotypic determinants: reagents for use with rat monoclonal antibodies. *Hybridoma*. 1982; 1(3):257-273.(Immunogen)

BD Biosciences

bdbiosciences.com

United States Canada Europe Japan Asia Pacific Latin America/Caribbean 877.232.8995 888.259.0187 32.53.720.550 0120.8555.90 65.6861.0633 55.11.5185.9995 For country-specific contact information, visit bdbiosciences.com/how to order/

Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton Dickinson and Company is strictly prohibited.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.

BD, BD Logo and all other trademarks are the property of Becton, Dickinson and Company. ©2006 BD

