

## Technical Data Sheet

## Biotin Mouse Anti-Rat IgG2b

## Product Information

Material Number:	553898
Size:	0.5 mg
Concentration:	0.5 mg/ml
Clone:	RG7/11.1
Immunogen:	Pooled rat IgG
Isotype:	Mouse (SJL) IgG2b, $\kappa$
Reactivity:	QC Testing: Rat
Storage Buffer:	Aqueous buffered solution containing $\leq 0.09\%$ sodium azide.

## Description

The RG7/11.1 antibody reacts specifically with the Fc region of rat IgG2b. It does not react with other Ig isotypes.

This antibody is routinely tested by flow cytometric analysis. Other applications were tested at BD Biosciences Pharmingen during antibody development only or reported in the literature.

## Preparation and Storage

The monoclonal antibody was purified from tissue culture supernatant or ascites by affinity chromatography.

The antibody was conjugated with biotin under optimum conditions, and unreacted biotin was removed.

Store undiluted at 4° C and protected from prolonged exposure to light. Do not freeze.

## Application Notes

## Application

Flow cytometry	Routinely Tested
ELISA Detection	Tested During Development

## Recommended Assay Procedure:

For the sandwich rat IgG2b ELISA, biotinylated mAb RG7/11.1 is optimal for detection with purified anti-rat IgG2b mAb G15-337 (Cat. No. 553882) for capture. Biotinylated RG7/11.1 mAb may be used as a secondary reagent in immunofluorescent staining. RG7/11.1 antibody is effective for detection of cell-surface or intracellular Ig by immunofluorescent staining with flow cytometric analysis. For flow cytometric detection of intracytoplasmic IgG2b, we recommend FITC-conjugated RG7/11.1 mAb (Cat. No 553900). For immunohistochemical staining, we recommend the use of biotinylated G15-337 mAb in our special formulation for immunohistochemistry, Cat. No. 550327.

## Suggested Companion Products

Catalog Number	Name	Size	Clone
553900	FITC Mouse Anti-Rat IgG2b	0.5 mg	RG7/11.1

## Product Notices

1. Since applications vary, each investigator should titrate the reagent to obtain optimal results.
2. Please refer to [wwwbdbiosciences.com/pharmingen/protocols](http://wwwbdbiosciences.com/pharmingen/protocols) for technical protocols.
3. Caution: Sodium azide yields highly toxic hydrazoic acid under acidic conditions. Dilute azide compounds in running water before discarding to avoid accumulation of potentially explosive deposits in plumbing.

## References

Springer TA, Bhattacharya A, Cardoza JT, Sanchez-Madrid F. Monoclonal antibodies specific for rat IgG1, IgG2a, and IgG2b subclasses, and kappa chain monotypic and allotypic determinants: reagents for use with rat monoclonal antibodies. *Hybridoma*. 1982; 1(3):257-273.(Immunogen)

## BD Biosciences

[wwwbdbiosciences.com](http://wwwbdbiosciences.com)

United States 877.232.8995    Canada 888.259.0187    Europe 32.53.720.550    Japan 0120.8555.90    Asia Pacific 65.6861.0633    Latin America/Caribbean 55.11.5185.9995

For country-specific contact information, visit [wwwbdbiosciences.com/how\\_to\\_order/](http://wwwbdbiosciences.com/how_to_order/)

Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton Dickinson and Company is strictly prohibited.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.

BD, BD Logo and all other trademarks are the property of Becton, Dickinson and Company. ©2007 BD

