

Technical Data Sheet

Biotin Hamster Anti-Mouse CD95**Product Information**


Material Number:	554256
Alternate Name:	Fas/APO-1
Size:	0.5 mg
Concentration:	0.5 mg/ml
Clone:	Jo2
Immunogen:	WR19L mouse lymphoma cells transformed with recombinant mouse Fas
Isotype:	Armenian Hamster IgG2, λ 2
Reactivity:	QC Testing: Mouse
Storage Buffer:	Aqueous buffered solution containing \leq 0.09% sodium azide.

Description

Fas antigen, CD95, is a 45 kDa cell-surface protein which can mediate apoptosis. It belongs to the TNF (tumor necrosis factor)/NGF receptor family. Expression of Fas has been described in the thymus, liver, heart, lung and ovary. Fas plays an important role in the apoptotic process that takes place during development. Monoclonal antibodies recognizing Fas such as Jo2 have cytolytic activity on cells expressing Fas. The cell death stimulated by Fas antibodies is characteristic of apoptosis and suggests that the lethal effects are a result of interaction of antibody with a functional Fas antigen as opposed to complement-mediated lysis.

The Jo2 antibody recognizes mouse Fas. The Jo2 antibody shows cytolytic activity against cell lines expressing mouse Fas by inducing apoptosis. Intraperitoneal injections of Jo2 mAb have been shown to kill mice and induce apoptotic hepatocyte death. Jo2 mAb immunoprecipitates mouse Fas as a 45 kDa band from W4 cells. W4 cells are WR19L mouse lymphoma cells transformed with mouse Fas. The difference between the observed MW of Fas and that deduced from its amino acid sequence (Mr 34,971) may be due to glycosylation.

This antibody is routinely tested by flow cytometric analysis. Other applications were tested at BD Biosciences Pharmingen during antibody development only or reported in the literature.



Expression of Fas antigen on mouse thymocytes analyzed by flow cytometry. Thymocytes from a BALB/c mouse were incubated with either Jo2 followed by a Streptavidin-PE conjugated second-step (Cat. No. 554061, filled histogram) or with only second step (open histogram). Jo2 specifically stained more than 90% of the cells.

Preparation and Storage

The monoclonal antibody was purified from tissue culture supernatant or ascites by affinity chromatography. The antibody was conjugated with biotin under optimum conditions, and unreacted biotin was removed. Store undiluted at 4° C and protected from prolonged exposure to light. Do not freeze.

Application Notes**Application**

Flow cytometry

Routinely Tested

BD Biosciences

bdbiosciences.com

United States	Canada	Europe	Japan	Asia Pacific	Latin America/Caribbean
877.232.8995	888.259.0187	32.53.720.550	0120.8555.90	65.6861.0633	55.11.5185.9995

For country-specific contact information, visit bdbiosciences.com/how_to_order/

Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton Dickinson and Company is strictly prohibited.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.

BD, BD Logo and all other trademarks are the property of Becton, Dickinson and Company. ©2006 BD

**BD Biosciences**

Suggested Companion Products

<u>Catalog Number</u>	<u>Name</u>	<u>Size</u>	<u>Clone</u>
554061	PE Streptavidin	0.5 mg	(none)

Product Notices

1. Although hamster immunoglobulin isotypes have not been well defined, BD Biosciences Pharmingen has grouped Armenian and Syrian hamster IgG monoclonal antibodies according to their reactivity with a panel of mouse anti-hamster IgG mAbs. A table of the hamster IgG groups, Reactivity of Mouse Anti-Hamster Ig mAbs, may be viewed at http://wwwbdbiosciences.com/pharmingen/hamster_chart_11x17.pdf.
2. Since applications vary, each investigator should titrate the reagent to obtain optimal results.
3. Please refer to wwwbdbiosciences.com/pharmingen/protocols for technical protocols.
4. Caution: Sodium azide yields highly toxic hydrazoic acid under acidic conditions. Dilute azide compounds in running water before discarding to avoid accumulation of potentially explosive deposits in plumbing.

References

- Enari M, Hug H, Nagata S. Involvement of an ICE-like protease in Fas-mediated apoptosis. *Nature*. 1995; 375(6526):78-81.(Clone-specific: Functional assay)
- Hiromatsu K, Aoki Y, Makino M, et al. Increased Fas antigen expression in murine retrovirus-induced immunodeficiency syndrome, MAIDS. *Eur J Immunol*. 1994; 24(10):2446-2451.(Clone-specific: Flow cytometry, Functional assay)
- Kagi D, Vignaux F, Ledermann B, et al. Fas and perforin pathways as major mechanisms of T cell-mediated cytotoxicity. *Science*. 1994; 265(5171):528-530. (Clone-specific: Flow cytometry, Functional assay, Immunoprecipitation)
- Nagata S. Apoptosis regulated by a death factor and its receptor: Fas ligand and Fas. *Philos Trans R Soc Lond B Biol Sci*. 1994; 345(1313):281-287. (Clone-specific: Functional assay)
- Nagata S. Fas and Fas ligand: a death factor and its receptor. *Adv Immunol*. 1994; 57:129-144.(Clone-specific: Functional assay)
- Ni R, Tomita Y, Matsuda K, et al. Fas-mediated apoptosis in primary cultured mouse hepatocytes. *Exp Cell Res*. 1994; 215(2):332-337.(Clone-specific: Functional assay)
- Ogasawara J, Suda T, Nagata S. Selective apoptosis of CD4+CD8+ thymocytes by the anti-Fas antibody. *J Exp Med*. 1995; 181(2):485-491.(Clone-specific: Flow cytometry, Functional assay)
- Ogasawara J, Watanabe-Fukunaga R, Adachi M, et al. Lethal effect of the anti-Fas antibody in mice. *Nature*. 1993; 364(6440):806-809.(Immunogen: Flow cytometry, Immunoprecipitation)
- Takahashi T, Tanaka M, Brannan CI, Jenkins NA, Copeland NG, Suda T, and Nagata S. Generalized lymphoproliferative disease in mice, caused by a point mutation in the Fas ligand. *Cell*. 1994; 76:969-976.(Biology)