

## Technical Data Sheet


## FITC Rat Anti-Human IL-2

## Product Information

|                  |                                                                  |
|------------------|------------------------------------------------------------------|
| Material Number: | 554565                                                           |
| Size:            | 0.1 mg                                                           |
| Concentration:   | 0.5 mg/ml                                                        |
| Clone:           | MQ1-17H12                                                        |
| Immunogen:       | Human IL-2 Recombinant Protein                                   |
| Isotype:         | Rat IgG2a, $\kappa$                                              |
| Reactivity:      | QC Testing: Human                                                |
| Storage Buffer:  | Aqueous buffered solution containing $\leq 0.09\%$ sodium azide. |

## Description

The MQ1-17H12 antibody reacts with human interleukin-2 (IL-2). The immunogen used to generate the MQ1-17H12 hybridoma was recombinant human IL-2. Unconjugated or purified forms of this antibody have been reported to be neutralizing for human IL-2 bioactivity.



**Expression of IL-2 by stimulated CD3<sup>+</sup> human PBMC.** Human PBMCs were stimulated for 6 hours with PMA (Sigma, Cat. #P-8139) and calcium ionophore A23187 (Sigma, Cat. #C-9275) in the presence of 2  $\mu$ M GolgiStop™ (Cat. No. 554724). The PBMCs were stained with PE-Cy™5-anti-CD3 (Cat. No. 555334), fixed, permeabilized, and then stained with 0.25  $\mu$ g of FITC-rat anti-human IL-2 antibody (Cat. No. 554565) (left panel). To demonstrate specificity of staining, the binding of FITC-MQ1-17H12 was blocked by the preincubation of the conjugated antibody with a molar excess of recombinant human IL-2 protein (1.0  $\mu$ g, Cat. No. 554603; middle panel), and by preincubation of the fixed/permeabilized cells with an excess of the unlabelled MQ1-17H12 antibody (10  $\mu$ g, Cat. No. 554563; right panel) prior to staining with the FITC-MQ1-17H12 antibody. The quadrant markers for the bivariate dot plots were set based on the autofluorescence control, and verified with the blocking controls.

## Preparation and Storage

Store undiluted at 4°C and protected from prolonged exposure to light. Do not freeze.

The monoclonal antibody was purified from tissue culture supernatant or ascites by affinity chromatography.

The antibody was conjugated with FITC under optimum conditions, and unreacted FITC was removed.

## Application Notes

## Application

|                                         |                  |
|-----------------------------------------|------------------|
| Intracellular staining (flow cytometry) | Routinely Tested |
|-----------------------------------------|------------------|

## Recommended Assay Procedure:

**Immunofluorescent Staining and Flow Cytometry:** The FITC-conjugated MQ1-17H12 antibody can be used for multicolor immunofluorescent staining and flow cytometric analysis to identify and enumerate IL-2-producing cells within mixed cell populations (see image). For optimal immunofluorescent staining for flow cytometric analysis, this anti-cytokine antibody should be titrated ( $\leq 0.5$   $\mu$ g mAb/ $1 \times 10^6$  cells). For specific methodology, please visit the protocol section on our web site, <http://www.bdbiosciences.com/resources/index.jsp>

A useful control for demonstrating specificity of staining is either of the following: 1) pre-block the conjugated MQ1-17H12 antibody with its ligand (e.g., recombinant human IL-2; Cat. No. 554603) prior to staining, or 2) pre-block the fixed/permeabilized cells with unlabelled MQ1-17H12 antibody (Cat. No. 554563) prior to staining. The intracellular cytokine staining technique and use of blocking controls are described in detail by C. Prussin and D. Metcalfe. A suitable rat IgG2a isotype control for assessing the level of background staining on paraformaldehyde-fixed/saponin-permeabilized mouse or human cells is FITC-R35-95 (Cat. No. 554688); use at comparable concentrations to the antibody of interest (e.g.,  $\leq 0.5$   $\mu$ g mAb/ $1 \times 10^6$  cells).

## BD Biosciences

[bdbiosciences.com](http://bdbiosciences.com)

| United States | Canada       | Europe        | Japan        | Asia Pacific | Latin America/Caribbean |
|---------------|--------------|---------------|--------------|--------------|-------------------------|
| 877.232.8995  | 800.979.9408 | 32.53.720.550 | 0120.8555.90 | 65.6861.0633 | 55.11.5185.9995         |

For country contact information, visit [bdbiosciences.com/contact](http://bdbiosciences.com/contact)

Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton, Dickinson and Company is strictly prohibited.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.

Unless otherwise noted, BD, BD Logo and all other trademarks are property of Becton, Dickinson and Company. © 2011 BD



**Neutralization/Blocking:** The NA/LE (Cat. No. 554562) format of the MQ1-17H12 antibody is useful for neutralization of human IL-2 bioactivity. A suitable NA/LE rat IgG2aisotype control is R35-95, Cat. No. 554687.

### Suggested Companion Products

| Catalog Number | Name                                              | Size      | Clone     |
|----------------|---------------------------------------------------|-----------|-----------|
| 554563         | Purified Rat Anti-Human IL-2                      | 0.1 mg    | MQ1-17H12 |
| 554603         | Recombinant Human IL-2                            | 10 µg     | (none)    |
| 554688         | FITC Rat IgG2a, κ Isotype Control                 | 0.1 mg    | R35-95    |
| 554724         | Protein Transport Inhibitor (Containing Monensin) | 0.7 ml    | (none)    |
| 555334         | PE-Cy™5 Mouse Anti-Human CD3                      | 100 tests | UCHT1     |
| 554562         | Purified NA/LE Rat Anti-Human IL-2                | 0.5 mg    | MQ1-17H12 |
| 554687         | Purified NA/LE Rat IgG2a κ Isotype Control        | 0.5 mg    | R35-95    |

### Product Notices

1. Since applications vary, each investigator should titrate the reagent to obtain optimal results.
2. Please refer to [www.bdbiosciences.com/pharming/en/protocols](http://www.bdbiosciences.com/pharming/en/protocols) for technical protocols.
3. Caution: Sodium azide yields highly toxic hydrazoic acid under acidic conditions. Dilute azide compounds in running water before discarding to avoid accumulation of potentially explosive deposits in plumbing.
4. This product is manufactured and sold under license from Pestka Biomedical Laboratories, Inc. (d/b/a PBL InterferonSource) and may be used solely as indicated. This product may not be resold or incorporated in any manner into another product for resale. Any use for therapeutics is strictly prohibited. This product is covered by U.S. Patent No. 5,597,901 and Bulgarian Patent No. BG1895.
5. Cy is a trademark of Amersham Biosciences Limited.
6. All other brands are trademarks of their respective owners.

### References

Abrams J. Immunoenzymetric assay of mouse and human cytokines using NIP-labeled anti-cytokine antibodies. In: Coligan J, Kruisbeek A, Margulies D, Shevach E, Strober W, ed. *Current Protocols in Immunology*. New York: John Wiley and Sons; 1995:6.20-6.21. (Biology)

Abrams JS, Roncarolo MG, Yssel H, Andersson U, Gleich GJ, Silver JE. Strategies of anti-cytokine monoclonal antibody development: immunoassay of IL-10 and IL-5 in clinical samples. *Immunol Rev*. 1992; 127:5-24. (Biology)

Prussin C, Metcalfe DD. Detection of intracytoplasmic cytokine using flow cytometry and directly conjugated anti-cytokine antibodies. *J Immunol Methods*. 1995; 188(1):117-128. (Clone-specific: Flow cytometry)

### BD Biosciences

[bdbiosciences.com](http://bdbiosciences.com)

|               |              |               |              |              |                         |
|---------------|--------------|---------------|--------------|--------------|-------------------------|
| United States | Canada       | Europe        | Japan        | Asia Pacific | Latin America/Caribbean |
| 877.232.8995  | 800.979.9408 | 32.53.720.550 | 0120.8555.90 | 65.6861.0633 | 55.11.5185.9995         |

For country contact information, visit [bdbiosciences.com/contact](http://bdbiosciences.com/contact)

Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton, Dickinson and Company is strictly prohibited.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.

Unless otherwise noted, BD, BD Logo and all other trademarks are property of Becton, Dickinson and Company. © 2011 BD

