Technical Data Sheet


PE Mouse Anti-Human CD8

Product Information

M 4 1 N 1	
Material Number:	555367
Alternate Name:	CD8α; CD8A; CD8 alpha; Leu2; MAL; T8; p32
Size:	100 Tests
Vol. per Test:	20 µl
Clone:	RPA-T8
Isotype:	Mouse IgG1, ĸ
Reactivity:	QC Testing: Human
	Tested in Development: Rhesus, Cynomolgus, Baboon
Workshop:	IV T171; V T-CD08.03; VI 6T-CD8.1, 6T-081
Storage Buffer:	Aqueous buffered solution containing BSA, protein stabilizer, and $\leq 0.09\%$ sodium azide.

Description

The RPA-T8 monoclonal antibody specifically binds to CD8 alpha (CD8a). CD8a is a type I transmembrane glycoprotein and a member of the immunoglobulin superfamily. CD8 α is expressed by the majority of thymocytes, by subpopulations of $\alpha\beta$ T cells and $\gamma\delta$ T cells and by some NK cells. Cell surface CD8a is expressed either as a disulfide-linked homodimer (CD8aa) or as a heterodimer (CD8aβ) when disulfide-bonded to a CD8 beta chain (CD8 β). CD8-positive $\alpha\beta$ T cells coexpress both CD8 $\alpha\alpha$ homodimers and CD8 $\alpha\beta$ heterodimers whereas some γδ T cells and NK cells express CD8αα homodimers. CD8 plays important roles in T cell activation and selection. The extracellular IgSF domain of CD8a binds to a non-polymorphic determinant on HLA class I molecules (a3 domain) and enables CD8 to function as a co-receptor with MHC class I-restricted TCR during T cell recognition of antigen. The cytoplasmic domain of CD8a associates with Lck, a Src family protein tyrosine kinase that is involved in intracellular signaling. The RPA-T8 and HIT8a monoclonal antibodies are not cross-blocking. This clone has been reported to react with a subset of peripheral blood lymphocytes, but not monocytes nor granuloyctes, of baboon and both rhesus and cynomolgus macaque monkey. In general, a higher frequency of CD8+ and CD4+CD8+ lymphocytes are observed in non-human primates compared to normal human donors.



Profile of peripheral blood lymphocytes analyzed on a FACScan (BDIS, San Jose, CA)

Preparation and Storage

Store undiluted at 4°C and protected from prolonged exposure to light. Do not freeze. The monoclonal antibody was purified from tissue culture supernatant or ascites by affinity chromatography. The antibody was conjugated with R-PE under optimum conditions, and unconjugated antibody and free PE were removed.

Application Notes

Application							
Flow cytom	etry				Routinely Tested		
Suggested	Companio	on Products	;				
Catalog Num	ber	Name				Size	Clone
555749		PE Mouse IgG1, κ Isotype Control				100 Tests	MOPC-21
BD Biosci							
bdbiosciences. United States		Europe	Japan	Asia Pacific	Latin America/Caribbean		S BI
877.232.8995	866.979.9408	32.2.400.98.95		65.6861.0633	55.11.5185.9995		
For country co	ntact informatio	on, visit bdbiosci	ences.com/conta	ct			
of any patents. B use of our produ product or as a co written authoriza	D Biosciences will n cts. Purchase does r omponent of anoth ation of Becton, Dic	ot be held responsil not include or carry a ner product. Any use kinson and Compar	ble for patent infring any right to resell or	gement or other vic transfer this produ er than the permitte d.	e the above product in violation lations that may occur with the ct either as a stand-alone ed use without the express		

Unless otherwise noted, BD, BD Logo and all other trademarks are property of Becton, Dickinson and Company. © 2015 BD

Product Notices

- 1. This reagent has been pre-diluted for use at the recommended Volume per Test. We typically use 1×10^{6} cells in a 100-µl experimental sample (a test).
- 2. An isotype control should be used at the same concentration as the antibody of interest.
- 3. Caution: Sodium azide yields highly toxic hydrazoic acid under acidic conditions. Dilute azide compounds in running water before discarding to avoid accumulation of potentially explosive deposits in plumbing.
- 4. Source of all serum proteins is from USDA inspected abattoirs located in the United States.
- 5. Species testing during development may have been performed with a different format of the same clone. Selected applications have been tested for cross-reactivity.
- 6. For fluorochrome spectra and suitable instrument settings, please refer to our Multicolor Flow Cytometry web page at www.bdbiosciences.com/colors.
- 7. Please refer to www.bdbiosciences.com/pharmingen/protocols for technical protocols.

References


Knapp W, Dörken B, Gilks WR, et al, ed. Leucocyte Typing IV. New York, NY: Oxford University Press; 1989:1-1182. (Biology) Schlossman SF, Boumsell L, Gilks W, et al, ed. Leukocyte Typing V: White Cell Differentiation Antigens. Oxford: Oxford University Press; 1995. (Clone-specific) Schlossman SF, Boumsell L, Gilks W, et al, ed. Leucocyte Typing V. New York: Oxford University Press; 1995. (Clone-specific)

BD Biosciences

bdbiosciences.com

United StatesCanadaEuropeJapanAsia Pacific877.232.8995866.979.940832.2.400.98.950120.8555.9065.6861.0633For country contact information, visit bdbiosciences.com/contact

Latin America/Caribbean 55.11.5185.9995



For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale. Unless otherwise noted, BD, BD Logo and all other trademarks are property of Becton, Dickinson and Company. © 2015 BD