
Technical Data Sheet PE Mouse Anti-Human CD33

Product Information	
Material Number:	555450
Size:	100 tests
Vol. per Test:	20 µl
Clone:	WM53
Isotype:	Mouse IgG1, ĸ
Reactivity:	QC Testing: Human
Workshop:	IV M505
Storage Buffer:	Aqueous buffered solution containing BSA and $\leq 0.09\%$ sodium azide.


Description

Reacts with a 67 kDa type I transmembrane glycoprotein expressed on monocytes, activated T cells, myeloid progenitors as well as mast cells. CD33 is absent on normal platelets, lymphocytes, erythrocytes, and hematopoietic stem cells. Reports indicate that this glycoprotein can function as a sialic acid-dependent cell adhesion molecule and this function can be modulated by endogenous sialoglycoconjugates when CD33 is expressed on the membrane.

This antibody is routinely tested by flow cytometric analysis. Other applications were tested at BD Biosciences Pharmingen during antibody development only or reported in the literature.



Profile of peripheral blood monocytes analyzed on a FACScan (BDIS, San Jose, CA)



Profile of peripheral blood granulocytes analyzed on a FACScan (BDIS, San Jose, CA)

Preparation and Storage

The monoclonal antibody was purified from tissue culture supernatant or ascites by affinity chromatography. The antibody was conjugated with R-PE under optimum conditions, and unconjugated antibody and free PE were removed by gel filtration chromatography. Store undiluted at 4° C and protected from prolonged exposure to light. Do not freeze.

Application Notes

Application

Flow cytometry				Routinely Tested	Routinely Tested	
BD Biosciences bdbiosciences.com						
United States 877.232.8995 For country-spe	Canada 888.259.0187 cific contact infor	Europe 32.53.720.550 mation, visit bdbio	Japan 0120.8555.90 sciences.com/how	Asia Pacific 65.6861.0633 _to_order/	Latin America/Caribbean 55.11.5185.9995	

Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton Dickinson and Company is stictly prohibited. For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale. BD, BD Logo and all other trademarks are the property of Becton, Dickinson and Company. ©2006 BD

BD Biosciences

Suggested Companion Products

Catalog Number	Name	Size	Clone
555749	PE Mouse IgG1 κ Isotype Control	100 tests	MOPC-21

Product Notices

- 1. This reagent has been pre-diluted for use at the recommended Volume per Test. We typically use 1 X 10e6 cells in a 100-µl experimental sample (a test).
- 2. Since applications vary, each investigator should titrate the reagent to obtain optimal results.
- 3. Please refer to www.bdbiosciences.com/pharmingen/protocols for technical protocols.
- 4. Caution: Sodium azide yields highly toxic hydrazoic acid under acidic conditions. Dilute azide compounds in running water before discarding to avoid accumulation of potentially explosive deposits in plumbing.
- 5. Source of all serum proteins is from USDA inspected abattoirs located in the United States.

References

Knapp W, Dorken B, et al, ed. Leucocyte Typing IV. New York: Oxford University Press; 1989.(Clone-specific)

Favaloro EJ, Bradstock KF, Kabral A, Grimsley P, Zowtyj H, Zola H. Further characterization of human myeloid antigens (gp160,95; gp150; gp67): investigation of epitopic heterogeneity and non-haemopoietic distribution using panels of monoclonal antibodies belonging to CD-11b, CD-13 and CD-33. *Br J Haematol.* 1988; 69(2):163-171.(Biology)

Favaloro EJ, Moraitis N, Koutts J, Exner T, Bradstock KF. Endothelial cells and normal circulating haemopoietic cells share a number of surface antigens. *Thromb* Haemost. 1989; 61(2):217-224.(Biology)

Freeman SD, Kelm S, Barber EK, Crocker PR. Characterization of CD33 as a new member of the sialoadhesin family of cellular interaction molecules. *Blood*. 1995; 85(8):2005-2012.(Biology)

Nakamura Y, Noma M, Kidokoro M, et al. Expression of CD33 antigen on normal human activated T lymphocytes. Blood. 1994; 83(5):1442-1443.(Biology)