Technical Data Sheet

Purified Mouse Anti-Human CD54

Product Information

 Material Number:
 555510

 Alternate Name:
 ICAM-1

 Size:
 0.1 mg

 Concentration:
 0.5 mg/ml

 Clone:
 HA58

 Isotype:
 Mouse IgG1 κ

Reactivity: QC Testing: Human


Workshop: VI A095

Storage Buffer: Aqueous buffered solution containing ≤0.09% sodium azide.

Description

Reacts with the 85-110 kDa integral membrane glycoprotein, also known as intracellular adhesion molecule-1 (ICAM-1), expressed on endothelial cells and both resting (weak) and activated (moderate) lymphocytes and monocytes. CD54 is a ligand for the leukocyte function antigen-1 (CD11a). CD54 antibodies are used for the studies of inflammatory processes and neoplasia. This antibody (NA/LE format) blocks the mixed-lymphocyte reaction (MLR) and the purified format is suitable for staining acetone-fixed, frozen tissue sections.

This antibody is routinely tested by flow cytometric analysis. Other applications were tested at BD Biosciences Pharmingen during antibody development only or reported in the literature.



Profile of resting peripheral blood lymphocytes analyzed on a FACScan (BDIS, San Jose, CA)

Preparation and Storage

The monoclonal antibody was purified from tissue culture supernatant or ascites by affinity chromatography. Store undiluted at 4° C.

Application Notes

Application

Flow cytometry Routinely Tested

BD Biosciences

bdbiosciences.com

United States Canada Europe Japan Asia Pacific Latin America/Caribbean 877.232.8995 888.259.0187 32.53.720.550 0120.8555.90 65.6861.0633 55.11.5185.9995 For country-specific contact information, visit bdbiosciences.com/how to order/

Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton Dickinson and Company is strictly prohibited.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.

BD, BD Logo and all other trademarks are the property of Becton, Dickinson and Company. ©2006 BD



Suggested Companion Products

Catalog Number	Name	Size	Clone	
555746	Purified Mouse IgG1 Kappa Isotype Control	0.1 mg	MOPC-21	
555988	FITC Goat Anti-Mouse IgG/IgM	0.5 mg	Gt/Ms	

Product Notices

- 1. Since applications vary, each investigator should titrate the reagent to obtain optimal results.
- 2. Please refer to www.bdbiosciences.com/pharmingen/protocols for technical protocols.
- Caution: Sodium azide yields highly toxic hydrazoic acid under acidic conditions. Dilute azide compounds in running water before
 discarding to avoid accumulation of potentially explosive deposits in plumbing.

References

Kishimoto T, von dem Borne AEG, Goyert SM,et al., ed. Leucocyte Typing VI: White Cell Differentiation Antigens. London: Garland Publishing; 1997. (Clone-specific)

Schlossman SF, Boumsell L, Gilks W, et al, ed. Leukocyte Typing V: White Cell Differentiation Antigens. New York: Oxford University Press; 1995.(Biology) Knapp W, Dorken B, et al, ed. Leucocyte Typing IV. New York: Oxford University Press; 1989.(Biology)

555510 Rev. 10 Page 2 of 2