Technical Data Sheet

Biotin Mouse Anti-Human CD56

Product Information

555515 **Material Number:** N-CAM Alternate Name: 100 tests Size: 20 µl Vol. per Test: B159 Clone:

Mouse IgG1, κ Isotype: QC Testing: Human Reactivity:

Workshop:


Storage Buffer: Aqueous buffered solution containing BSA, protein stabilizer, and ≤0.09%

sodium azide.

Description

Reacts with the 220/135 kDa heavily glycosylated antigen isoforms present on a subpopulation of peripheral blood large granular lymphocytes which demonstrate natural killer activity, but not on myeloid cells, erythrocytes or B cells. This antigen is a pan-NK-cell marker. CD56 is virtually identical to an isoform of the neutral cell adhesion molecule (NCAM), a structure mediating homotypic and heterotypic cell-cell interactions.

This antibody is routinely tested by flow cyotmetric analysis. Other applications were tested at BD Biosciences Pharmingen during antibody development only or reported in the literature.



Profile of peripheral blood lymphocytes analyzed on a FACScan (BDIS, San Jose, CA)

Preparation and Storage

The monoclonal antibody was purified from tissue culture supernatant or ascites by affinity chromatography. The antibody was conjugated with biotin under optimum conditions, and unreacted biotin was removed.

Store undiluted at 4° C and protected from prolonged exposure to light. Do not freeze.

Application Notes

Application

Flow cytometry Routinely Tested

BD Biosciences

bdbiosciences.com

United States Europe Japan Asia Pacific Latin America/Caribbean 888.259.0187 32.53.720.550 0120.8555.90 877.232.8995 65.6861.0633 55.11.5185.9995

For country-specific contact information, visit bdbiosciences.com/how_to_order/

Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton Dickinson and Company is strictly prohibited.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale. BD, BD Logo and all other trademarks are the property of Becton, Dickinson and Company. ©2006 BD



Suggested Companion Products

Catalog Number	Name	Size	Clone
554061	Streptavidin PE	0.5 mg	(none)
555747	Biotin Mouse IgG1 κ Isotype Control	100 tests	MOPC-21

Product Notices

- 1. This reagent has been pre-diluted for use at the recommended Volume per Test. We typically use 1 X 10e6 cells in a 100-μl experimental sample (a test).
- 2. Since applications vary, each investigator should titrate the reagent to obtain optimal results.
- 3. Please refer to www.bdbiosciences.com/pharmingen/protocols for technical protocols.
- Caution: Sodium azide yields highly toxic hydrazoic acid under acidic conditions. Dilute azide compounds in running water before discarding to avoid accumulation of potentially explosive deposits in plumbing.
- 5. Source of all serum proteins is from USDA inspected abattoirs located in the United States.

References

Schlossman SF, Boumsell L, Gilks W, et al, ed. *Leukocyte Typing V: White Cell Differentiation Antigens*. New York: Oxford University Press; 1995.(Clone-specific) Schlossman S, Boumell L, et al, ed. *Leucocyte Typing V*. New York: Oxford University Press; 1995.(Biology) Guesdon JL, Ternynck T, Avrameas S. The use of avidin-biotin interaction in immunoenzymatic techniques. *J Histochem Cytochem*. 1979; 27(8):1131-1139. (Biology)

555515 Rev. 9 Page 2 of 2