

## Technical Data Sheet

## Purified Mouse Anti-Human CD64


## Product Information

Material Number:	555525
Size:	0.1 mg
Concentration:	0.5 mg/ml
Clone:	10.1
Isotype:	Mouse IgG1, $\kappa$
Reactivity:	QC Testing: Human
Workshop:	VI MA36
Storage Buffer:	Aqueous buffered solution containing $\leq 0.09\%$ sodium azide.

## Description

The monoclonal antibody 10.1 reacts with CD64, a 75 kD type I transmembrane glycoprotein that is a high affinity receptor for IgG (Fc $\gamma$ RI). CD64, a key receptor in the development of immune responses, has a dual role as a low affinity receptor for IgG3 and high affinity receptor for IgG2a linking innate and adaptive immunities. CD64 mediates endocytosis, phagocytosis, antibody-dependent cellular toxicity, cytokine release and superoxide generation. CD64 is expressed largely on macrophages and dendritic cells.

This antibody is routinely tested by flow cytometric analysis. Other applications were tested at BD Biosciences Pharmingen during antibody development only or reported in the literature.



Profile of peripheral blood monocytes analyzed on a FACSscan (BDIS, San Jose, CA)

## Preparation and Storage

The monoclonal antibody was purified from tissue culture supernatant or ascites by affinity chromatography.

Store undiluted at 4°C.

## Application Notes

## Application

Flow cytometry	Routinely Tested
Immunohistochemistry-frozen	Tested During Development

## Suggested Companion Products

Catalog Number	Name	Size	Clone
555746	Purified Mouse IgG1, $\kappa$ Isotype Control	0.1 mg	MOPC-21
555988	FITC Goat Anti-Mouse IgG/IgM	0.5 mg	Polyclonal

## Product Notices

- Since applications vary, each investigator should titrate the reagent to obtain optimal results.

## BD Biosciences

bdbiosciences.com

United States	Canada	Europe	Japan	Asia Pacific	Latin America/Caribbean
877.232.8995	888.259.0187	32.53.720.550	0120.8555.90	65.6861.0633	55.11.5185.9995

For country-specific contact information, visit [bdbiosciences.com/how\\_to\\_order/](http://bdbiosciences.com/how_to_order/)

Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton Dickinson and Company is strictly prohibited.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.

BD, BD Logo and all other trademarks are the property of Becton, Dickinson and Company. ©2008 BD



2. Please refer to [www.bdbiosciences.com/pharmingen/protocols](http://www.bdbiosciences.com/pharmingen/protocols) for technical protocols.
3. Caution: Sodium azide yields highly toxic hydrazoic acid under acidic conditions. Dilute azide compounds in running water before discarding to avoid accumulation of potentially explosive deposits in plumbing.

## References

- Dougherty GJ, Selvendran Y, Murdoch S, Palmer DG, Hogg N. The human mononuclear phagocyte high-affinity Fc receptor, FcRI, defined by a monoclonal antibody, 10.1. *Eur J Immunol.* 1987; 17(10):1453-1459. (Biology)
- Indik ZK, Hunter S, Huang MM, et al. The high affinity Fc gamma receptor (CD64) induces phagocytosis in the absence of its cytoplasmic domain: the gamma subunit of Fc gamma RIIIA imparts phagocytic function to Fc gamma RI. *Exp Hematol.* 1994; 22(7):599-606. (Biology)
- Kishimoto T, von dem Borne AEG, Goyert SM, et al., ed. *Leucocyte Typing VI: White Cell Differentiation Antigens*. London: Garland Publishing; 1997. (Clone-specific)
- Schlossman S, Boumell L, et al, ed. *Leucocyte Typing V*. New York: Oxford University Press; 1995. (Biology)