


Technical Data Sheet

Biotin Mouse Anti-Human CD80**Product Information**

Material Number:	555682
Alternate Name:	B7-1
Size:	100 tests
Vol. per Test:	20 µl
Clone:	BB1
Isotype:	Mouse IgM, κ
Reactivity:	QC Testing: Human
Storage Buffer:	Aqueous buffered solution containing BSA and ≤0.09% sodium azide.

Description

Reacts with a 60 kD transmembrane glycoprotein which was clustered as CD80 in the Fifth International Workshop on Human Differentiation Antigens. There have been several recent studies reporting discrepancies in the expression and function of CD80 when the BB1 clone is used compared to other anti-CD80 monoclonals (e.g., L307.4). Caution should therefore be used in the interpretation of data generated using BB1, as reports indicate that BB1 may recognize, in addition to CD80, CD74 and even possibly another as yet undefined molecule. CD80, a member of the Ig supergene family, is expressed on activated B cells, macrophages, and dendritic cells. It is the ligand for two molecules expressed on T cells: CD28 and CD152 (CTLA-4). CD80 is also expressed in activated CD4+ and CD8+ T cells, appearing late after activation, suggesting that activated T cells may be capable of autocrine costimulation via the CD28 activation pathway. The binding of CD28 molecule by CD28 mAb or by CD80 antigen results in T-cell activation and a signal for IL-2 production.



*Profile of RAJI cells
analyzed by flow cytometry*

Preparation and Storage

The monoclonal antibody was purified from tissue culture supernatant or ascites by affinity chromatography.

The antibody was conjugated with biotin under optimum conditions, and unreacted biotin was removed.

Store undiluted at 4° C and protected from prolonged exposure to light. Do not freeze.

Application Notes**Application**

Flow cytometry

Routinely Tested

BD Biosciences
www.bdbiosciences.com

United States	Canada	Europe	Japan	Asia Pacific	Latin America/Caribbean
877.232.8995	888.259.0187	32.53.720.550	0120.8555.90	65.6861.0633	55.11.5185.9995

For country-specific contact information, visit www.bdbiosciences.com/how_to_order/

Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton Dickinson and Company is strictly prohibited.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.

BD, BD Logo and all other trademarks are the property of Becton, Dickinson and Company. ©2007 BD



Suggested Companion Products

Catalog Number	Name	Size	Clone
554061	PE Streptavidin	0.5 mg	(none)
555582	Biotin Mouse IgM κ Isotype Control	100 tests	G155-228

Product Notices

1. This reagent has been pre-diluted for use at the recommended Volume per Test. We typically use 1 X 10⁶ cells in a 100- μ l experimental sample (a test).
2. Since applications vary, each investigator should titrate the reagent to obtain optimal results.
3. Please refer to www.bdbiosciences.com/pharmlingen/protocols for technical protocols.
4. Caution: Sodium azide yields highly toxic hydrazoic acid under acidic conditions. Dilute azide compounds in running water before discarding to avoid accumulation of potentially explosive deposits in plumbing.
5. Source of all serum proteins is from USDA inspected abattoirs located in the United States.

References

- Schlossman SF, Bousmell L, Gilks W, et al, ed. *Leukocyte Typing V: White Cell Differentiation Antigens*. New York: Oxford University Press; 1995. (Clone-specific)
- Azuma M, Yssel H, Phillips JH, Spits H, Lanier LL. Functional expression of B7/BB1 on activated T lymphocytes. *J Exp Med*. 1993; 177(3):845-850. (Clone-specific)
- Behrens L, Kerschensteiner M, Misgeld T, Goebels N, Wekerle H, Hohlfield R. Human muscle cells express a functional costimulatory molecule distinct from B7.1 (CD80) and B7.2 (CD86) in vitro and in inflammatory lesions. *J Immunol*. 1998; 161(11):5943-5951.(Clone-specific)
- Freeman GJ, Cardoso AA, Boussiotis VA, et al. The BB1 monoclonal antibody recognizes both cell surface CD74 (MHC class II-associated invariant chain) as well as B7-1 (CD80), resolving the question regarding a third CD28/CTLA-4 counterreceptor. *J Immunol*. 1998; 161(6):2708-2715.(Clone-specific)
- Guesdon JL, Ternynck T, Avrameas S. The use of avidin-biotin interaction in immunoenzymatic techniques. *J Histochem Cytochem*. 1979; 27(8):1131-1139. (Methodology)
- Koulova L, Clark EA, Shu G, Dupont B. The CD28 ligand B7/BB1 provides costimulatory signal for alloactivation of CD4+ T cells. *J Exp Med*. 1991; 173(3):759-762.(Clone-specific)
- Schwartz RH. Costimulation of T lymphocytes: the role of CD28, CTLA-4, and B7/BB1 in interleukin-2 production and immunotherapy. *Cell*. 1992; 71(7):1065-1068.(Clone-specific)