# **Technical Data Sheet**

# Biotin Mouse IgG1 κ Isotype Control

#### **Product Information**

 Material Number:
 555747

 Size:
 100 tests

 Vol. per Test:
 20 μl

 Clone:
 MOPC-21


**Isotype:** Mouse (BALB/c) IgG1 κ

Storage Buffer: Aqueous buffered solution containing BSA and ≤0.09% sodium azide.

#### Description

The MOPC-21 immunoglobulin is a mouse myeloma protein. The MOPC-21 immunoglobulin was selected as an isotype control following screening for low background on a variety of mouse and human tissues.

This antibody is routinely tested by flow cytometric analysis. Other applications were tested at BD Biosciences Pharmingen during antibody development only or reported in the literature.



Profile of peripheral blood lymphocytes analyzed by flow cytometry.

# **Preparation and Storage**

The monoclonal antibody was purified from tissue culture supernatant or ascites by affinity chromatography. The antibody was conjugated with biotin under optimum conditions, and unreacted biotin was removed. Store undiluted at 4° C and protected from prolonged exposure to light. Do not freeze.

### **Application Notes**

#### Application

Immunocytochemistry (cytospins)	Tested During Development
Flow cytometry	Routinely Tested

#### **Product Notices**

- Caution: Sodium azide yields highly toxic hydrazoic acid under acidic conditions. Dilute azide compounds in running water before
  discarding to avoid accumulation of potentially explosive deposits in plumbing.
- 2. Source of all serum proteins is from USDA inspected abattoirs located in the United States.
- 3. Since applications vary, each investigator should titrate the reagent to obtain optimal results.
- This reagent has been pre-diluted for use at the recommended Volume per Test. We typically use 1 X 10e6 cells in a 100-μl experimental sample (a test).
- 5. Please refer to www.bdbiosciences.com/pharmingen/protocols for technical protocols.

### **BD** Biosciences

bdbiosciences.com

United States Canada Europe Japan Asia Pacific Latin America/Caribbean 877.232.8995 888.259.0187 32.53.720.550 0120.8555.90 65.6861.0633 55.11.5185.9995 For country-specific contact information, visit bdbiosciences.com/how to order/

Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton Dickinson and Company is strictly prohibited.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.

BD, BD Logo and all other trademarks are the property of Becton, Dickinson and Company, ©2006 BD



### References

Guesdon JL, Ternynck T, Avrameas S. The use of avidin-biotin interaction in immunoenzymatic techniques. *J Histochem Cytochem*. 1979; 27(8):1131-1139. (Biology)

555747 Rev. 7 Page 2 of 2