

Technical Data Sheet


Purified Mouse Anti-Human CD79a

Product Information

Material Number:	555934
Alternate Name:	Igα, mb-1
Size:	0.1 mg
Concentration:	0.5 mg/ml
Clone:	HM47
Isotype:	Mouse IgG1, κ
Reactivity:	QC Testing: Human Cross Reactivity Confirmed in Development: Mouse
Workshop:	V cB017
Storage Buffer:	Aqueous buffered solution containing ≤0.09% sodium azide.

Description

Reacts with the cytoplasmic domain of a 47 kDa membrane glycoprotein present on B lymphocytes, also referred to as mb-1. It is expressed on B cells at various stages of differentiation, from pre-B cell stage, probably before expression of cytoplasmic μ chain, to plasma cell stage, in which it is detected only in the cytoplasm. CD79a associates with CD79b to form part of the B-cell receptor complex. It has been suggested that CD79a may play a role in mediating the transport of IgM to the cell surface. This antibody has been found to react on permeabilized A20 cells (mouse B cell line).



Profile of permeabilized peripheral blood lymphocytes with Cytofix/Cytoperm (Cat# 554714) analyzed on a FACScan (BDIS, San Jose, CA)

Preparation and Storage

The monoclonal antibody was purified from tissue culture supernatant or ascites by affinity chromatography.

Store undiluted at 4° C.

Application Notes

Application

Intracellular staining (flow cytometry)	Routinely Tested
---	------------------

Suggested Companion Products

Catalog Number	Name	Size	Clone
555746	Purified Mouse IgG1, κ Isotype Control	0.1 mg	MOPC-21
555988	FITC Goat Anti-Mouse IgG/IgM	0.5 mg	Polyclonal
554714	BD Cytofix/Cytoperm Fixation/Permeabilization Kit	250 tests	(none)

BD Biosciences

www.bdbiosciences.com

United States 877.232.8995 Canada 888.259.0187 Europe 32.53.720.550 Japan 0120.8555.90 Asia Pacific 65.6861.0633 Latin America/Caribbean 55.11.5185.9995

For country-specific contact information, visit www.bdbiosciences.com/how_to_order/

Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton Dickinson and Company is strictly prohibited.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.

BD, BD Logo and all other trademarks are the property of Becton, Dickinson and Company. ©2007 BD



Product Notices

1. Since applications vary, each investigator should titrate the reagent to obtain optimal results.
2. Please refer to www.bdbiosciences.com/pharming/en/protocols for technical protocols.
3. Caution: Sodium azide yields highly toxic hydrazoic acid under acidic conditions. Dilute azide compounds in running water before discarding to avoid accumulation of potentially explosive deposits in plumbing.

References

Schlossman SF, Boumsell L, Gilks W, et al, ed. *Leukocyte Typing V: White Cell Differentiation Antigens*. New York: Oxford University Press; 1995.(Clone-specific)
Mason DY, Cordell JL, Tse AG, et al. The IgM-associated protein mb-1 as a marker of normal and neoplastic B cells. *J Immunol*. 1991; 147(11):2474-2482.
(Biology)
Sakaguchi N, Kashiwamura S, Kimoto M, Thalmann P, Melchers F. B lymphocyte lineage-restricted expression of mb-1, a gene with CD3-like structural properties.
EMBO J. 1988; 7(11):3457-3464.(Biology)