# Technical Data Sheet Purified Mouse Anti-Human CD3ε

# **Product Information**

Material Number:	556610
Size:	0.1 mg
Concentration:	0.5 mg/ml
Clone:	SP34
Isotype:	Mouse IgG3, $\lambda$
Reactivity:	Human
	QC Testing: Baboon or Cynomolgus or Rhesus
Workshop:	V 5T-CD03.05
Storage Buffer:	Aqueous buffered solution containing protein stabilizer and $\leq 0.09\%$ sodium azide.

## Description

Clone SP34 reacts with human form of the 20 kDa epsilon subunit of the CD3 antigen/T-cell receptor (TCR) complex. This clone also cross-reacts with a major subset of peripheral blood lymphocytes, but not monocytes or granulocytes, of rhesus or cynomolgus and pigtail macaque monkeys. The distribution on lymphocytes is similar to that observed with normal human donor lymphocytes, with the majority of CD3-positive cells being negative when dual stained with antibodies to B or NK cell markers.

This antibody is routinely tested by flow cytometric analysis. Other applications were tested at BD Biosciences Pharmingen during antibody development only or reported in the literature.





#### **Preparation and Storage**

The monoclonal antibody was purified from tissue culture supernatant or ascites by affinity chromatography. Store undiluted at 4°C.

#### Application Notes

	•	
Ap	plication	

Flow cytometry	Routinely Tested

# Suggested Companion Products

Catalog Number	Name	Size	Clone
556657	Purified Mouse IgG3, ĸ Isotype Control	0.1 mg	J606
555988	FITC Goat Anti-Mouse IgG/IgM	0.5 mg	Polyclonal

#### **Product Notices**

1. Since applications vary, each investigator should titrate the reagent to obtain optimal results.

2. Please refer to www.bdbiosciences.com/pharmingen/protocols for technical protocols.

### **BD Biosciences**

bdbiosciences.com								
United States	Canada	Europe	Japan	Asia Pacific	Latin America/Caribbean			
877.232.8995	888.259.0187	32.53.720.550	0120.8555.90	65.6861.0633	55.11.5185.9995			
For country-specific contact information, visit bdbiosciences.com/how_to_order/								
Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton Dickinson and Company is strictly prohibited. For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale. BD, BD Logo and all other trademarks are the property of Becton, Dickinson and Company. @2008 BD								



 Caution: Sodium azide yields highly toxic hydrazoic acid under acidic conditions. Dilute azide compounds in running water before discarding to avoid accumulation of potentially explosive deposits in plumbing.

#### References

Sancho J, Ledbetter JA, Choi MS, Kanner SB, Deans JP, Terhorst C. CD3-zeta surface expression is required for CD4-p56lck-mediated upregulation of T cell antigen receptor-CD3 signaling in T cells. J Biol Chem. 1992; 267(11):7871-7879.(Biology) Schlossman SF, Boumsell L, Gilks W, et al, ed. Leukocyte Typing V: White Cell Differentiation Antigens. New York: Oxford University Press; 1995.(Clone-specific)