Technical Data Sheet Alexa Fluor® 700 Hamster IgG2, λ Isotype Control

Product Information	
Material Number:	557985
Alternate Name:	Anti-KLH
Size:	0.1 mg
Concentration:	0.2 mg/ml
Clone:	Ha4/8
Immunogen:	Keyhole limpet hemocyanin
Isotype:	Armenian Hamster IgG2, $\lambda 1$
Storage Buffer:	Aqueous buffered solution containing protein stabilizer and $\leq 0.09\%$ sodium azide.

Description

The Ha4/8 antibody is specific for the keyhole limpet hempcyanin (KLH).

Preparation and Storage

The monoclonal antibody was purified from tissue culture supernatant or ascites by affinity chromatography. The antibody was conjugated to Alexa Fluor® 700 under optimum conditions, and unreacted Alexa Fluor® 700 was removed. Store undiluted at 4° C and protected from prolonged exposure to light. Do not freeze.

Application Notes

Application	
Flow cytometry	Routinely Tested

Recommended Assay Procedure:

An isotype control should be used at the same concentration as the antibody of interest.

Note: Alexa Fluor® 700 conjugates of the hamster IgG2 isotype reportedly can non-specifically stain myeloid cells from mouse bone marrow.

Product Notices

- 1. Please refer to www.bdbiosciences.com/pharmingen/protocols for technical protocols.
- 2. Alexa Fluor® 700 has an adsorption maximum of ~700nm and a peak fluorescence emission of ~720nm. Before staining cells with this reagent, please confirm that your flow cytometer is capable of exciting the fluorochrome and discriminating the resulting fluorescence.
- 3. For fluorochrome spectra and suitable instrument settings, please refer to our Fluorochrome Web Page at www.bdbiosciences.com/pharmingen/colors.
- 4. Although hamster immunoglobulin isotypes have not been well defined, BD Biosciences Pharmingen has grouped Armenian and Syrian hamster IgG monoclonal antibodies according to their reactivity with a panel of mouse anti-hamster IgG mAbs. A table of the hamster IgG groups, Reactivity of Mouse Anti-Hamster Ig mAbs, may be viewed at http://www.bdbiosciences.com/pharmingen/hamster_chart_11x17.pdf.
- Caution: Sodium azide yields highly toxic hydrazoic acid under acidic conditions. Dilute azide compounds in running water before 5. discarding to avoid accumulation of potentially explosive deposits in plumbing.
- 6. The Alexa Fluor®, Pacific BlueTM, and Cascade Blue® dye antibody conjugates in this product are sold under license from Molecular Probes, Inc. for research use only, excluding use in combination with microarrays, or as analyte specific reagents. The Alexa Fluor® dyes (except for Alexa Fluor® 430), Pacific Blue™ dye, and Cascade Blue® dye are covered by pending and issued patents.

References

Mendrick DL, Kelly DM. Temporal expression of VLA-2 and modulation of its ligand specificity by rat glomerular epithelial cells in vitro. Lab Invest. 1993; 69(6):690-702.(Immunogen)

BD Biosciences

www.bdbiosciences.com				
atin America/Caribbean				
5.11.5185.9995				
For country-specific contact information, visit www.bdbiosciences.com/how_to_order /				
Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton Dickinson and Company is strictly prohlibited. For Research Use Only Net for use in diagnostic or theraneutic procedures. Not for resale				
5. de or				