BD[™] Cytometric Bead Array (CBA)



Technical Data Sheet Rat IFN-y Flex Set

Product Information

Material Number:
Size:
Bead Position:
Assay Range:
Reactivity:

558305 100 Tests A6 40-10,000 pg/mL QC Testing: Rat

Component Description: Component Mat. No: Component Storage Buffer:

Component Description: Component Mat. No: Component Storage Buffer:

Component Storage Buffer:

sodium azide. Component Description: Component Mat. No: 51-9005242

Rat IFN-y Standard 51-9004111 Lyophilized in an aqueous buffered solution containing BSA and ProClin™ 150.

Rat IFN-y PE Detection Reagent 51-9004156 Aqueous buffered solution containing BSA and ≤0.09%

Rat IFN-y Capture Bead A6 Aqueous buffered solution containing fetal bovine serum and $\leq 0.09\%$ sodium azide.




Figure 1. Example BD CBA Rat IFN-y Flex Set standard curve. Data acquired on a BD FACSArray bioanalyzer and analyzed using FCAP Array Software.

Description

The BDTM CBA Rat IFN-y Flex Set is a bead-based immunoassay capable of measuring rat interferon-y (IFN-y) in serum and cell culture supernatant samples. Rat reactivity was determined by testing samples with the BD CBA Rat IFN-y Flex Set. The biology and function of IFN-y has been extensively reviewed in the literature. For more information on bead-based immunoassays, refer to the product insert for the BD CBA Mouse/Rat Soluble Protein Master Buffer Kit (Cat. No. 558266 or 558267).

Preparation and Storage

This BDTM CBA Flex Set contains one vial each of Capture Bead and PE Detection Reagent and two vials of Standard. The Capture Bead and PE Detection Reagent components of this flex set have been formulated to a 50x concentration to ensure product performance when multiplexed. The Standard component is lyophilized and should be transferred to a 15 mL polypropylene tube for reconstitution. When reconstituted in 4.0 mL Assay Diluent, the standard has a protein concentration of 10,000 pg/mL. Discard unused reconstituted standard, do not store or reuse. Store lyophilized standard and other components at 4°C. Protect Capture Beads and the PE Detection Reagent from prolonged exposure to light.

Application Notes

Recommended Assay Procedure: The BD CBA Rat IFN-y Flex Set must be used in conjunction with a BD CBA Mouse/Rat Soluble Protein Master Buffer Kit (Cat. No. 558266, 100 tests, or 558267, 500 tests), a flow cytometer, and FCAP Array™ Software. Detailed instructions on the use of this product can be found in the manual for the BD CBA Mouse/Rat Soluble Protein Master Buffer Kit. When following the directions in the Master Buffer Kit, the top standard point for the BD CBA Rat IFN-y Flex Set will be 10,000 pg/mL. An example standard curve is shown in figure 1.

BD Biosciences

bdbiosciences.	om				
United States	Canada	Europe	Japan	Asia Pacific	Latin America/Caribbean
877.232.8995	800.268.5430	32.2.400.98.95	0120.8555.90	65.6861.0633	55.11.5185.9995
For country cor	ntact informatio	on, visit bdbioscie	ences.com/conta	ct	

Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton, Dickinson and Company is stictly prohibited. For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale

Unless otherwise noted, BD, BD Logo and all other trademarks are property of Becton, Dickinson and Company. © 2014 BD

When multiplexing the BD CBA Rat IFN- γ Flex Set assay with the BD CBA Rat IL-4 Flex Set assay (Cat. No. 558307), significantly higher background will be seen in the rat IL-4 assay. This increased background, while reducing the sensitivity of the rat IL-4 assay, will not effect the quantitation of rat IFN- γ or IL-4 in any other way.

The BD CBA Rat IFN-γ Flex Set should not be used in the same assay well with any non-BD CBA Rat Soluble Protein Flex Set reagents (such as BD CBA Human or Mouse Soluble Protein or Cell Signaling Flex Sets). For an updated assay compatibility chart for the BD CBA Rat Soluble Protein Flex Sets, please refer to the BD CBA Flex Set System homepage at http://www.bdbiosciences.com/cbasetup.

Performance

Limit of Detection: The theoretical limit of detection is 6.8 pg/mL and was determined by evaluating the estimated result of the average MFI of the negative control (0 pg/mL, n=30) + 2 standard deviations.

Specificity		Inter-Assay Reproducibility			Intra-Assay Reproducibility		
		Mean (pg/ml)	Standard Deviation	%CV	Mean (pg/ml)	Standard Deviation	%CV
	Sample 1	154.2	12.6	8%	139.7	2.5	2%
Rat IFN-γ	Sample 2	611.5	16.6	3%	601.1	9.9	2%
	Sample 3	2448.2	77.1	3%	2556.3	61.1	2%

Reproducibility: The inter-assay and intra-assay reproducibility were determined for the BD CBA Rat IFN-y Flex Set by evaluating ten replicates of three different sample levels (inter-assay) and two replicates of three different sample levels from four separate experiments (intra-assay) respectively.

		ulture natant	Serum		
Sample Dilution	Average % Recovery	Range	Average % Recovery	Range	
Rat IFN-γ	70%	68 - 74%	62%	60 - 67%	

Recovery: Cell culture supernatant and pooled rat serum were spiked with three different levels of protein. The spiked samples were assayed and the results were compared with expected values. Serum samples were diluted 1:4 before the protein was spiked into each. Serum used was commercially available pooled rat serum.

Comple Dilution		culture natant	Serum		
Sample Dilution	Detected (pg/ml)	% of Expected	Detected (pg/ml)	% of Expected	
Spiked sample	1987.8	100%	1859.7	100%	
1:2	989.4	100%	952.2	102%	
1:4	502.7	101%	489.0	105%	

Linearity: Cell culture supernatant and 1:4 diluted pooled rat serum were spiked with protein and serially diluted. The diluted samples were assayed and the results were compared with the original spiked sample.

Product Notices

1. ProClin is a trademark of Rohm and Haas Company.

with local/regional/national/international regulations.

- 2. Source of all serum proteins is from USDA inspected abattoirs located in the United States.
- 3. Caution: Sodium azide yields highly toxic hydrazoic acid under acidic conditions. Dilute azide compounds in running water before discarding to avoid accumulation of potentially explosive deposits in plumbing.

 Warning: CBA lyophilized standard contains 0.02% (w/w) of a CMIT/MIT mixture (3:1), which is a mixture of: 5-chloro-2-methyl-4-isothiazolin-3-one [EC No 247-500-7] and 2-methyl-4-isothiazolin-3-one [EC No 220-239-6] (3:1). Hazard statement: May cause an allergic skin reaction. Precautionary statements: Wear protective gloves/eye protection. Wear protective clothing. Avoid breathing mist/vapours/spray. If skin irritation or rash occurs: Get medical advice/attention. IF ON SKIN: Wash with plenty of water. Dispose of contents/container in accordance

BD Biosciences

bdbiosciences.	com				
United States 877.232.8995	Canada 800.268.5430	Europe 32.2.400.98.95	Japan 0120.8555.90	Asia Pacific 65.6861.0633	Latin America/Caribbean 55.11.5185.9995
For country co	ntact informatio	on, visit bdbioscie	ences.com/conta	ct	
of any patents. BL use of our produc product or as a co	D Biosciences will n ts. Purchase does n mponent of anoth tion of Becton, Dic	ot be held responsik Iot include or carry a er product. Any use kinson and Compan	ble for patent infrin any right to resell or of this product oth y is stictly prohibite	gement or other vio r transfer this produ er than the permitte d.	e the above product in violation lations that may occur with the ct either as a stand-alone ed use without the express

Unless otherwise noted, BD, BD Logo and all other trademarks are property of Becton, Dickinson and Company. © 2014 BD