

Technical Data Sheet

PE Mouse anti-PLC- γ 2 (pY759)


Product Information

Material Number:	558490
Size:	50 tests
Vol. per Test:	20 μ l
Clone:	K86-689.37
Immunogen:	Phosphorylated Human PLC- γ 2 Peptide
Isotype:	Mouse IgG1, κ
Reactivity:	Confirmed: Human Predicted: Mouse, Rat
Storage Buffer:	Aqueous buffered solution containing BSA and $\leq 0.09\%$ sodium azide.

Description

The Phospholipase C (PLC) isozymes hydrolyze phosphatidyl inositol biphosphate to inositol triphosphate and diacylglycerol. The former causes release of calcium from the endoplasmic reticulum, while the latter is an activator of Protein Kinase C. Within the PLC family, PLC- γ is the only member that contains SH2 and SH3 domains. These domains enable it to interact with protein tyrosine kinases and become enzymatically activated via phosphorylation. It exists as two isoforms: PLC- γ 1, which is ubiquitously expressed, and PLC- γ 2, which is found primarily in the hematopoietic system. PLC- γ 2-null mice exhibit impaired B lymphocyte maturation and defects in Fc receptor functions. Phosphorylation of PLC- γ 2 at tyrosines 753 and 759 (Y759) is required for activation of PLC- γ 2 enzyme activity. PLC- γ 2 phosphorylation at Y759 can be induced by stimulation of the B cell receptor in Ramos cells, the collagen receptor in platelets, and the T cell receptor in Jurkat cells, and it occurs downstream of Btk and BLNK in the signaling cascade of activated B cells.

The K86-689.37.73 antibody recognizes PLC- γ 2 phosphorylated at Y759 in the SH2-SH3 linker region.



Analysis of PLC- γ 2 (pY759) in human Burkitt's lymphoma. Ramos cells were either stimulated (open histogram) by cross-linking of surface IgM with purified Goat anti-Human IgM (SouthernBiotech) at 37°C for 3 minutes or unstimulated (solid histogram). The cells were fixed with pre-warmed BD™ Phosflow Fix Buffer I (Cat. No. 557870) for 10 minutes at 37°C, then permeabilized with BD™ Phosflow Perm Buffer III (Cat. No. 558050) on ice for at least 10 minutes, and then stained with PE Mouse anti-PLC- γ 2 (pY759). Flow cytometry was performed on a BD™ FACSCalibur flow cytometry system.

Preparation and Storage

The antibody was conjugated with R-PE under optimum conditions, and unconjugated antibody and free PE were removed. Store undiluted at 4°C and protected from prolonged exposure to light. Do not freeze.

Application Notes

Application

Intracellular staining (flow cytometry)	Tested
---	--------

Suggested Companion Products

Catalog Number	Name	Size	Clone
557870	Fix Buffer I	250 ml	(none)
558050	Perm Buffer III	125 ml	(none)

BD Biosciences

bdbiosciences.com

United States	Canada	Europe	Japan	Asia Pacific	Latin America/Caribbean
877.232.8995	800.979.9408	32.53.720.550	0120.8555.90	65.6861.0633	55.11.5185.9995

For country contact information, visit bdbiosciences.com/contact

Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton, Dickinson and Company is strictly prohibited.
For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.
Unless otherwise noted, BD, BD Logo and all other trademarks are property of Becton, Dickinson and Company. © 2011 BD



Product Notices

1. Please refer to www.bdbiosciences.com/pharming/en/protocols for technical protocols.
2. This reagent has been pre-diluted for use at the recommended Volume per Test. We typically use 1×10^6 cells in a 100- μ l experimental sample (a test).
3. Caution: Sodium azide yields highly toxic hydrazoic acid under acidic conditions. Dilute azide compounds in running water before discarding to avoid accumulation of potentially explosive deposits in plumbing.
4. Source of all serum proteins is from USDA inspected abattoirs located in the United States.

References

Kim YJ, Sekiya F, Poulin B, Bae YS, Rhee SG. Mechanism of B-cell receptor-induced phosphorylation and activation of phospholipase C- γ 2. *Mol Cell Biol.* 2004; 24(22):9986-9999. (Biology)

Ozdener F, Dangelmaier C, Ashby B, Kunapuli SP, Daniel JL. Activation of phospholipase C γ 2 by tyrosine phosphorylation.. *Mol Pharmacol.* 2002; 62(3):672-679. (Biology)

Wang D, Feng J, Wen R, et al. Phospholipase C γ 2 is essential in the functions of B cell and several Fc receptors. *Immunity.* 2000; 13:25-35. (Biology)

Wen R, Jou S-T, Chen Y, Hoffmeyer A, Wang D. Phospholipase C γ 2 is essential for specific functions of Fc ϵ R and Fc γ R. *J Immunol.* 2002; 169:6743-6752. (Biology)

BD Biosciences

bdbiosciences.com

United States	Canada	Europe	Japan	Asia Pacific	Latin America/Caribbean
877.232.8995	800.979.9408	32.53.720.550	0120.8555.90	65.6861.0633	55.11.5185.9995

For country contact information, visit bdbiosciences.com/contact

Conditions: The information disclosed herein is not to be constructed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton, Dickinson and Company is strictly prohibited.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.

Unless otherwise noted, BD, BD Logo and all other trademarks are property of Becton, Dickinson and Company. © 2011 BD

