Technical Data Sheet Alexa Fluor® 488 Mouse anti-LAT (pY171)

Product Information

Material Number:	558519
Size:	50 tests
Vol. per Test:	20 µl
Clone:	I58-1169
Immunogen:	Phosphorylated Human LAT
Isotype:	Mouse (BALB/c) IgG1, ĸ
Reactivity:	Human
Storage Buffer:	Aqueous buffered solution containing BSA and ≤0.09% sodium azide.

Description

Engagement of the T cell receptor (TCR) induces signal transduction pathways that enhance gene transcription and cellular proliferation and differentiation. TCR ligation results in the recruitment and activation of multiple protein tyrosine kinases (PTKs), including lck, fyn, and ZAP70. Adaptor proteins, such as Grb2 and SLP-76, relay the signal to downstream effector molecules. LAT (linker for activation of T cells) is a substrate of the activated ZAP70 and functions to bridge the activated TCR and its associated PTKs with tyrosine kinase substrates. LAT is expressed as 36- and 38-kDa forms that result from post-translational modification, and as a 42-kDa form that results from alternative splicing. LAT is an integral membrane protein that is phosphorylated at five tyrosine sites upon TCR ligation. Following phosphorylation, LAT binds a number of important signaling molecules, including Grb2, Vav, PLCγ1, and the p85 subunit of PI3K. Multiple studies have shown that functional LAT is required for T lymphocyte activation and thymocyte development.

The I58-1169 monoclonal antibody recognizes the phosphorylated tyrosine 171 (pY171) of LAT, which is one of the phosphotyrosine sites required for binding phosphoinositide 3-kinase, Grb2, and Gads.



Analysis of LAT (pY171) in human peripheral blood lymphocytes. Human peripheral blood mononuclear cells were either stimulated by cross-linking of CD3 and CD28 with NA/LE Mouse anti-Human CD3 mAb UCHT1 (Cat. No. 555329) and NA/LE Mouse anti-Human CD28 mAb CD28.2 (Cat. No. 555725) on ice for 15 minutes followed by Purified Goat anti-Mouse Ig (Cat. No. 553998) on ice for 15 minutes, and then allowed to undergo phosphorylation at 37°C for 1-2 minutes (shaded histogram) or unstimulated (open histogram). The cells were fixed (BD Cytofix™ buffer, Cat. No. 554655) for 10 minutes at 37 ℃, permeabilized (BD™ Phosflow Perm Buffer III. Cat. No. 558050) on ice for 30 minutes, blocked with normal mouse immunoglobulin, and then stained with Alexa Fluor® 488 Mouse anti-LAT (pY171) and PE Mouse anti-human CD3 mAb UCHT1 (Cat. No. 555333). The figure displays the upregulated phosphorylation of LAT in activated CD3-positive T lymphocytes. Flow cytometry was performed on a BD™ FACSCalibur flow cytometry system.

Preparation and Storage

The antibody was conjugated to Alexa Fluor® 488 under optimum conditions, and unreacted Alexa Fluor® 488 was removed. Store undiluted at 4°C and protected from prolonged exposure to light. Do not freeze.

Application Notes

Application	l .							
Intracellular staining (flow cytometry)				Routinely Tested				
Suggeste	d Compani	on Product	ts					
Catalog Nu	mber	Name				Size	Clone	
558050		Perm Bu	iffer III			125 ml	(none)	
554655		Fixation	Buffer			100 ml	(none)	
Product N 1. Please	Notices refer to www	bdbiosciences.	.com/pharmin	gen/protocols f	for technical protocols.			
BD Biosci	ences							
bdbiosciences	.com							
United States 877.232.8995	Canada 888.259.0187	Europe 32.53.720.550	Japan 0120.8555.90	Asia Pacific 65.6861.0633	Latin America/Caribbean 55.11.5185.9995			

 United States
 Canada
 Europe
 Japan
 Asia Pacific
 Latin America/Caribbean

 877.232.8995
 888.259.0187
 32.53.720.550
 0120.8555.90
 65.6861.0633
 55.11.5185.9995

 For country-specific contact information, visit
 bdbiosciences.com/how_to_order/
 Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product on a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton Dickinson and Company is strictly prohibited.
 For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.

 BD, BD Logo and all other trademarks are the property of Becton, Dickinson and Company. ©2008 BD
 Becton Biotecon Strictly and the trademarks are the property of Becton, Bickinson and Company.
 S2008 BD



- 2. This reagent has been pre-diluted for use at the recommended Volume per Test. We typically use 1×10^{6} cells in a 100-µl experimental sample (a test).
- 3. The Alexa Fluor®, Pacific Blue™, and Cascade Blue® dye antibody conjugates in this product are sold under license from Molecular Probes, Inc. for research use only, excluding use in combination with microarrays, or as analyte specific reagents. The Alexa Fluor® dyes (except for Alexa Fluor® 430), Pacific Blue™ dye, and Cascade Blue® dye are covered by pending and issued patents.
- 4. Alexa Fluor® 488 fluorochrome emission is collected at the same instrument settings as for fluorescein isothiocyanate (FITC).
- Caution: Sodium azide yields highly toxic hydrazoic acid under acidic conditions. Dilute azide compounds in running water before discarding to avoid accumulation of potentially explosive deposits in plumbing.
- 6. Source of all serum proteins is from USDA inspected abattoirs located in the United States.
- 7. For fluorochrome spectra and suitable instrument settings, please refer to our Fluorochrome Web Page at www.bdbiosciences.com/colors.
- 8. Alexa Fluor® is a registered trademark of Life Technologies Corporation.

References

Cho S, Velikovsky CA, Swaminathan CP, Houtman JCD, Samelson LE, Mariuzza RA. Structural basis for differential recognition of tyrosine-phosphorylated sites in the linker for activation of T cells (LAT) by the adaptor Gads. *EMBO J.* 2004; 23:1441-1451. (Biology)

Janssen E, Zhang W. Adaptor proteins in lymphocyte activation. Curr Opin Immunol. 2003; 15:269-276. (Biology)

Lin J, Weiss A. Identification of the minimal tyrosine residues required for linker for activation of T cell function. J Biol Chem. 2001; 276(31):29588-29595. (Biology) Paz PE, Wang S, Clarke H, Lu X, Stokoe D, Abo A. Mapping the ZAP-70 phosphorylation sites on LAT (linker for activation of T cells) required for recruitment and activation of signalling proteins in T cells. Biochem J. 2001; 356:461-471. (Biology)

Samelson LE. Signal transduction mediated by the T cell antigen receptor: The role of adapter proteins. *Annu Rev Immunol.* 2002; 20:371-394. (Biology) Zhu M, Janssen E, Zhang W. Minimal requirement of tyrosine residues of linker for activation of T cells in TCR signaling and thymocyte development. *J Immunol.* 2003; 170:325-333. (Biology)