

## Technical Data Sheet


## APC Rat anti-Mouse CD25

## Product Information

<b>Material Number:</b>	<b>558643</b>
<b>Alternate Name:</b>	IL-2 Receptor $\alpha$ chain, p55
<b>Size:</b>	0.1 mg
<b>Concentration:</b>	0.2 mg/ml
<b>Clone:</b>	3C7
<b>Isotype:</b>	Rat (LEW) IgG2b, $\kappa$
<b>Reactivity:</b>	Mouse
<b>Storage Buffer:</b>	Aqueous buffered solution containing $\leq 0.09\%$ sodium azide.

## Description

The 3C7 antibody reacts with CD25, the low affinity IL-2 Receptor (IL-2R $\alpha$ , p55) expressed on activated T and B lymphocytes from all mouse strains tested. IL-2R $\alpha$  by itself is not a signaling receptor. However, it can combine with IL-2 Receptor  $\beta$  (CD122) and  $\gamma_c$  (CD132) chains to form high-affinity, signaling receptor complexes for IL-2. Resting T and B lymphocytes and resting and activated NK cells do not express IL-2R $\alpha$ . CD25 is transiently expressed at a low level during normal B-cell development in the bone marrow on the CD45R/B220low TdT- sIg- Pre-B/Pre-B-II and CD45R/B220low TdT- sIgM+ sIgD- immature B stages, but not on the CD45R/B220low TdT+ sIg- Pro-B/Pre B-I stage nor on CD45R/B220high TdTIgM+ sIgD+ mature B cells. It is expressed at a higher level during a very early stage of T-cell development in fetal and adult thymus. Peripheral CD25+ CD4+ T lymphocytes called regulatory T (Treg) cells are involved in the maintenance of self-tolerance. It has also been reported that dendritic cells express CD25, recognized by mAb 7D4 (Cat. No. 553068). The 3C7 antibody recognizes an epitope of CD25 which is distinct from those recognized by mAbs 7D4 and PC61 (Cat. No. 553866), and it blocks binding of IL-2 to CD25.



**Flow cytometric analysis of APC-conjugated anti-mouse CD25 on mouse splenocytes.**

Concanavlin A activated murine splenocytes were stained with either APC anti-CD25(clone 3C7, Cat. No. 558643, solid line) or an APC rat IgG2b isotype control (catalog number 553991, dashed line) and analyzed by flow cytometry. Flow cytometry was performed on a BD FACSCalibur™ System and the histograms were derived from the gated events based on light scattering characteristics of viable splenocytes.

## Preparation and Storage

The polyclonal antibody was purified from antiserum by affinity chromatography.

The antibody was conjugated to APC under optimum conditions, and unconjugated antibody and free APC were removed.

Store undiluted at 4°C and protected from prolonged exposure to light. Do not freeze.

## Application Notes

## Application

Flow cytometry	Routinely Tested
----------------	------------------

## Product Notices

1. Please refer to [www.bdbiosciences.com/pharmingen/protocols](http://www.bdbiosciences.com/pharmingen/protocols) for technical protocols.
2. Caution: Sodium azide yields highly toxic hydrazoic acid under acidic conditions. Dilute azide compounds in running water before discarding to avoid accumulation of potentially explosive deposits in plumbing.
3. For fluorochrome spectra and suitable instrument settings, please refer to our Fluorochrome Web Page at [www.bdbiosciences.com/colors](http://www.bdbiosciences.com/colors).

## References

## BD Biosciences

[bdbiosciences.com](http://bdbiosciences.com)

United States	Canada	Europe	Japan	Asia Pacific	Latin America/Caribbean
877.232.8995	888.268.5430	32.53.720.550	0120.8555.90	65.6861.0633	0800.771.7157

For country-specific contact information, visit [bdbiosciences.com/how\\_to\\_order/](http://bdbiosciences.com/how_to_order/)

Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton Dickinson and Company is strictly prohibited.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.

BD, BD Logo and all other trademarks are the property of Becton, Dickinson and Company. ©2011 BD



Chen J, Ma A, Young F, Alt FW. IL-2 receptor alpha chain expression during early B lymphocyte differentiation. *Int Immunol*. 1994; 6(8):1265-1268. (Biology)

Crowley M, Inaba K, Witmer-Pack M, Steinman RM. The cell surface of mouse dendritic cells: FACS analyses of dendritic cells from different tissues including thymus. *Cell Immunol*. 1989; 118(1):108-125. (Biology)

Habu S, Okumura K, Diamantstein T, Shevach EM. Expression of interleukin 2 receptor on murine fetal thymocytes. *Eur J Immunol*. 1985; 15(5):456-460. (Clone-specific)

Malek TR, Schmidt JA, Shevach EM. The murine IL 2 receptor. III. Cellular requirements for the induction of IL 2 receptor expression on T cell subpopulations. *J Immunol*. 1985; 134(4):2405-2413. (Biology)

Moreau JL, Nabholz M, Diamantstein T, Malek T, Shevach E, Theze J. Monoclonal antibodies identify three epitope clusters on the mouse p55 subunit of the interleukin 2 receptor: relationship to the interleukin 2-binding site. *Eur J Immunol*. 1987; 17(7):929-935. (Biology)

Ortega G, Robb RJ, Shevach EM, Malek TR. The murine IL 2 receptor. I. Monoclonal antibodies that define distinct functional epitopes on activated T cells and react with activated B cells. *J Immunol*. 1984; 133(4):1970-1975. (Immunogen)

Pollard AM, Lipscomb MF. Characterization of murine lung dendritic cells: similarities to Langerhans cells and thymic dendritic cells. *J Exp Med*. 1990; 172(1):159-167. (Biology: Controls)

Read S, Malmstrom V, Powrie F. Cytotoxic T lymphocyte-associated antigen 4 plays an essential role in the function of CD25(+)CD4(+) regulatory cells that control intestinal inflammation. *J Exp Med*. 2000; 192(2):295-302. (Biology)

Rolink A, Grawunder U, Winkler TH, Karasuyama H, Melchers F. IL-2 receptor alpha chain (CD25, TAC) expression defines a crucial stage in pre-B cell development. *Int Immunol*. 1994; 6(8):1257-1264. (Biology)

Taniguchi T, Minami Y. The IL-2/IL-2 receptor system: a current overview. *Cell*. 1993; 73(1):5-8. (Biology)