Technical Data Sheet

Purified Mouse Anti-Human CD135

Product Information


558995 **Material Number:** 0.1 mg Size: 0.5 mg/ml **Concentration:** 4G8 Clone: Mouse IgG1, κ Isotype:

QC Testing: Human Reactivity:

Storage Buffer: Aqueous buffered solution containing ≤0.09% sodium azide.

Description

Reacts with FMS-like tyrosine kinase 3 (flt3), a 155-160 kD, class III tyrosine kinase receptor (CD135), which is structurally related to the receptors for PDGF, SCF-1 and kit ligand. CD135 is expressed on multipotential, myelomonocytic and primitive B-cell progenitors. The most primitive hematopoietic progenitor cells express low levels of CD135. CD135 plays the role of growth factor receptor for early hematopoietic progenitors. Reports from studies with knockout mice have shown that targeted CD135 disruption impaired the development of primitive progenitor cells of all hematopoietic lineages with significant impact on lymphopoietic precursors.



Profile of REH cell line analyzed by flow cytometry

Preparation and Storage

The monoclonal antibody was purified from tissue culture supernatant or ascites by affinity chromatography. Store undiluted at 4° C.

Application Notes

Application

- 1	TI		
- 1	Flow cytometry	Routinely Tested	
- 1	1 low cytometry	Routinery residu	

Suggested Companion Products

Catalog Number	Name	Size	Clone
555746	Purified Mouse IgG1, κ Isotype Control	0.1 mg	MOPC-21
555988	FITC Goat Anti-Mouse IgG/IgM	0.5 mg	Polyclonal

BD Biosciences

www.bdbiosciences.com

Europe 32.53.720.550 0120.8555.90 877.232.8995 888.259.0187 65.6861.0633 55.11.5185.9995 For country-specific contact information, visit www.bdbiosciences.com/how_to_order/

Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation Conditions: The information disclosed nerein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton Dickinson and Company is strictly prohibited.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.

BD, BD Logo and all other trademarks are the property of Becton, Dickinson and Company. ©2007 BD



Product Notices

- 1. Since applications vary, each investigator should titrate the reagent to obtain optimal results.
- 2. Please refer to www.bdbiosciences.com/pharmingen/protocols for technical protocols.
- 3. Caution: Sodium azide yields highly toxic hydrazoic acid under acidic conditions. Dilute azide compounds in running water before discarding to avoid accumulation of potentially explosive deposits in plumbing.

References

Kishimoto T, von dem Borne AEG, Goyert SM,et al., ed. Leucocyte Typing VI: White Cell Differentiation Antigens. London: Garland Publishing; 1997. (Clone-specific)

Lyman SD, James L, Johnson L. Cloning of the human homologue of the murine flt3 ligand: a growth factor for early hematopoietic progenitor cells. *Blood.* 1994; 83(10):2795-27801.(Biology)

Mackarehtschian K, Hardin JD, Moore KA, Boast S, Goff SP, Lemischka IR. Targeted disruption of the flk2/flt3 gene leads to deficiencies in primitive hematopoietic progenitors. *Immunity*. 1995; 3(1):147-161.(Biology)

pold I, Ziegler BL, Köhler I. Functional and phenotypic characterization of cord blood and bone marrow subsets expressing FLT3 (CD135) receptor tyrosine kinase. Blood. 1997; 90(1):111-125.(Biology)

Rosnet O, Schiff C, Pébusque MJ. Human FLT3/FLK2 gene: cDNA cloning and expression in hematopoietic cells. *Blood.* 1993; 82(4):1110-1119.(Biology) Rusten LS, Lyman SD, Veiby OP, Jacobsen SE. The FLT3 ligand is a direct and potent stimulator of the growth of primitive and committed human CD34+ bone marrow progenitor cells in vitro. *Blood.* 1996; 87(4):1317-1325.(Biology)

558995 Rev. 4 Page 2 of 2