

Technical Data Sheet

Purified Mouse Anti-Human TNF for ICC


Product Information

Material Number:	559071
Size:	0.25 mg
Concentration:	0.5 mg/ml
Clone:	MAb11
Immunogen:	Recombinant Human TNF
Isotype:	Mouse IgG1, κ
Reactivity:	QC Testing: Human
Storage Buffer:	Aqueous buffered solution containing $\leq 0.09\%$ sodium azide.

Description

The MAb11 antibody reacts with human tumor necrosis factor (TNF, also known as TNF- α) protein. TNF is an efficient paracrine and endocrine mediator of inflammatory and immune functions. It regulates the growth and differentiation of a variety of cell types. TNF is cytotoxic for transformed cells when in conjunction with IFN- γ . It is secreted by activated monocytes/macrophages and other cells such as B cells, T cells and fibroblasts. The immunogen used to generate the MAb11 hybridoma was recombinant human TNF. The use of the MAb11 antibody has been reported to cross-react with TNF of rhesus monkey.

This antibody is routinely tested by immunocytochemical staining. Other applications were tested at BD Biosciences Pharmingen during antibody development only or reported in the literature.



PBMC were isolated from human peripheral blood by density gradient centrifugation and were cultured for 2 days with plate bound anti-human CD3 and soluble anti-human CD28 in the presence of human recombinant IL-2 and human recombinant IL-4. The cells were subsequently harvested, washed and recultured with human recombinant IL-2 and human recombinant IL-4 for an additional 3 days. Finally, the cells were harvested, washed and stimulated with PMA (Sigma, Cat. No. P-8139, 5 ng/ml) and ionomycin (Sigma, Cat. No. I-0634, 500 ng/ml) in the presence of GolgiStop™ (Cat. No. 554724) for 4 hr at 37°C. The activated cells were harvested and the presence of TNF producing cells was detected by immunocytochemistry using a three-step staining procedure that employs a Biotin Goat anti-Mouse IgG secondary antibody (Cat. No. 550337) and a horseradish peroxidase-based detection system (Streptavidin-HRP Cat. No. 550947) (Nomarski optics, original magnification 400X). To demonstrate specificity of staining the binding of the MAb11 (Cat. No. 559071) antibody was blocked by the preincubation of the purified antibody with excess recombinant human TNF protein (Cat. No. 554589; data not shown).

Preparation and Storage

The monoclonal antibody was purified from tissue culture supernatant or ascites by affinity chromatography. Store undiluted at 4° C.

Application Notes

Application

Immunocytochemistry (cytospins)	Routinely Tested
---------------------------------	------------------

Recommended Assay Procedure:

Immunocytochemistry: The ICC format of the purified MAb11 (Cat. No. 559071) antibody can be used to identify and enumerate human TNF producing cells by immunocytochemistry. For optimal indirect immunocytochemical staining, the MAb11 antibody should be titrated and visualized via a three-step staining procedure. Please see protocol below for a detailed description of our recommended procedure.

BD Biosciences

bdbiosciences.com

United States	Canada	Europe	Japan	Asia Pacific	Latin America/Caribbean
877.232.8995	888.259.0187	32.53.720.550	0120.8555.90	65.6861.0633	55.11.5185.9995

For country-specific contact information, visit bdbiosciences.com/how_to_order/

Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton Dickinson and Company is strictly prohibited.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.

BD, BD Logo and all other trademarks are the property of Becton, Dickinson and Company. ©2006 BD



BD

BD Biosciences

CYTOKINE IMMUNOCYTOCHEMISTRY PROTOCOL

REAGENTS REQUIRED

1. Fixation Buffer: 5% formalin (10% formalin, CMS, Cat. No. 245-684) is dissolved in phosphate buffered-saline (PBS) (Bacto FA Buffer, Difco Laboratories, Cat. No. 2314-15-0), or BD Pharmingen™ ICC Fixation Buffer (BD Cat. No. 550010)
2. Endogenous Peroxidase Blocking Buffer: DAKO Peroxidase Blocking Reagent (DAKO, Cat. No. S2001).
3. Endogenous Biotin Blocking Buffer: Biotin/Avidin Blocking Kit (Vector Laboratories, Cat. No. SP-2001).
4. Antibody dilution buffer: BD Pharmingen™ IHC Antibody Diluent Buffer supplemented with saponin (Cat. No. 559148).
5. Microscopic slides: Adhesion Slides (Erie Scientific Company, Cat. No. ER-202B-AD) or for cytopins, Colorfrost /Plus slides (Fisher, Cat. No. 12-550-17).
6. Detection system: BD Pharmingen™ Streptavidin-horseradish peroxidase (HRP), (Cat. No. 550946), or Anti-Mouse Ig HRP Detection Kit (Cat. No. 551011).
7. Mounting medium for short-term storage: Aqua-mount® (Lerner Laboratories, Cat. No. 13800).
8. DAB Substrate Kit (contains 3-3'-Diaminobenzidine tetra hydrochloride), (BD Cat. No. 550880), or Anti-Mouse Ig HRP Detection Kit (Cat. No. 551011).

SECONDARY ANTIBODIES

Biotin Goat anti-Mouse IgG (Cat. No. 550337) or Anti-Mouse Ig HRP Detection Kit (Cat. No. 551011)

PROCEDURE FOR IMMUNOCYTOCHEMICAL STAINING OF SINGLE-CELL PREPARATIONS

This procedure describes the immunoenzymatic technique of staining cytokines within individual cells that are immobilized on microscopic slides via adherence (adherent slides) or centrifugation (cytopins).

ADHESION SLIDES

1. Harvest cells and wash them twice in PBS using centrifugation (400 x g for 5 min) to remove residual protein.
2. Adjust the cell concentration at 4-5 x 10⁶ cells/ml in PBS.
3. Place 20 µl of the cell suspension in each well of the adhesion slides and let them adhere at room temperature (RT) for 20 min. Please note that the slides should be washed in PBS at RT for 5 min before transferring the cells.
4. Fix cells on slides using fixation buffer for 15 min at RT.
5. Wash slides 2X in PBS with 5 min incubations.
6. Block slides with PBS supplemented with 1% (w/v) BSA (Sigma, Cat. No. A43-78) for 30 min at RT or 10 min at 37°C.
7. Wash slides 2X in PBS and proceed with staining or air dry them and store them at -80°C for future use.
8. Incubate slides with 20 µl of 1% goat serum and PBS with 0.1% (w/v) saponin for 30 min at RT.
9. Wash slides 2X with PBS with 5 min incubations.
10. Block endogenous peroxidase activity with Endogenous Peroxidase Blocking Buffer (20 µl/well) for 10 min at RT.
11. Wash 2X in PBS with 5 min incubations.
12. Incubate each well with Avidin (20 µl/well) for 15 min.
13. Wash 2X in PBS with 5 min incubations.
14. Incubate each well with Biotin (20 µl/well) for 15 min.
15. Wash 2X in PBS with 5 min incubations.
16. Incubate each well for 1 hr at RT with 20 µl of purified cytokine-specific antibody or appropriate immunoglobulin isotype control diluted in IHC Diluent Buffer (Cat. No. 559148) supplemented with saponin.
17. Wash slides 2X in PBS with 5 min incubations.
18. Incubate each well with 20 µl of a biotinylated secondary antibody diluted in IHC Diluent Buffer for 30 min at RT.
19. Wash 2X in PBS with 5 min incubations.
20. Apply 20 µl of Streptavidin-HRP (BD Cat. No. 550946) to each well on slides and incubate for 30 min at RT.
21. Wash slides 2X with PBS with 5 minutes incubations.
22. Incubate with DAB Substrate as directed, (BD Cat. No. 550880) for less than 5 min at RT.
23. Stop the development of the color reaction by washing with PBS.
24. The slides are subsequently mounted in short-term storage mounting medium.

CYTOSPINS

1. Assemble the Cytospin's sample chamber (e.g. Cytospin 3, Shandon, UK or comparable centrifuge), filter card, slide and cytopsin racks according to manufacturer's specifications.
2. Load 40 µl of approximately 1 x 10⁶ cells to each sample chamber.
3. Spin slides at 600 rpm for 2 min.
4. Take slides out of the cytopsin rack and place them on a staining rack.
5. For fixation and staining please follow the steps 4 through 24 specified above for staining cells on adhesion slides.

Suggested Companion Products

Catalog Number	Name	Size	Clone
550010	ICC Fixation Buffer	100 ml	(none)
559148	Antibody Diluent for IHC	125 ml	(none)
551011	Anti-Mouse Ig HRP Detection Kit	200 tests	(none)
554724	Protein Transport Inhibitor (Containing Monensin)	0.7 ml	(none)
550880	DAB Substrate Kit	500 tests	(none)
550946	Streptavidin HRP	50 ml	(none)
550337	Biotin Goat Anti-Mouse Igs	1.0 ml	Polyclonal

Product Notices

1. Since applications vary, each investigator should titrate the reagent to obtain optimal results.
2. Please refer to www.bdbiosciences.com/pharmlingen/protocols for technical protocols.
3. Caution: Sodium azide yields highly toxic hydrazoic acid under acidic conditions. Dilute azide compounds in running water before discarding to avoid accumulation of potentially explosive deposits in plumbing.

References

Danis VA, Franic GM, Rathjen DA, Brooks PM. Effects of granulocyte-macrophage colony-stimulating factor (GM-CSF), IL-2, interferon-gamma (IFN-gamma), tumour necrosis factor-alpha (TNF-alpha) and IL-6 on the production of immunoreactive IL-1 and TNF-alpha by human monocytes. *Clin Exp Immunol*. 1991; 85(1):143-150.(Clone-specific: ELISA)

Hsu SM, Raine L, Fanger H. Use of avidin-biotin-peroxidase complex (ABC) in immunoperoxidase techniques: a comparison between ABC and unlabeled antibody (PAP) procedures. *J Histochem Cytochem*. 1981; 29(4):577-580.(Methodology: Immunocytochemistry (cytospins))

Hsu SM, Raine L, Fanger H. A comparative study of the peroxidase-antiperoxidase method and an avidin-biotin complex method for studying polypeptide hormones with radioimmunoassay antibodies. *Am J Clin Pathol*. 1981; 75(5):734-738.(Methodology: Immunocytochemistry (cytospins))

Rathjen DA, Cowan K, Furphy LJ, Aston R. Antigenic structure of human tumour necrosis factor: recognition of distinct regions of TNF alpha by different tumour cell receptors. *Mol Immunol*. 1991; 28(1-2):79-86.(Clone-specific: ELISA)