

## Technical Data Sheet

## Purified Mouse Anti-Rat CD106

## Product Information

<b>Material Number:</b>	<b>559165</b>
<b>Alternate Name:</b>	VCAM-1
<b>Size:</b>	0.5 mg
<b>Concentration:</b>	0.5 mg/ml
<b>Clone:</b>	MR106
<b>Immunogen:</b>	Rat VCAM-1-transfected L5178Y Cells
<b>Isotype:</b>	Mouse IgG1, $\kappa$
<b>Reactivity:</b>	QC Testing: Rat
<b>Storage Buffer:</b>	Aqueous buffered solution containing $\leq 0.09\%$ sodium azide.

## Description

The MR106 antibody reacts with CD106, the ~110 kDa transmembrane vascular cell adhesion molecule-1 (VCAM-1). Expression of CD106 on endothelial cells is induced by inflammatory cytokines and in certain pathologic conditions. In the rat bone marrow, CD106 has been reported to be expressed on dendritic cells and subsets of the myeloid lineage, but not detectable on peripheral leukocytes. CD106 has been reported to be a counter-receptor for VLA-4 ( $\alpha 4\beta 1$  integrin) and  $\alpha 4\beta 7$  integrin, and it plays an important role in leukocyte extravasation, particularly at sites of inflammation.

This antibody is routinely tested by flow cytometric analysis. Other applications were tested at BD Biosciences Pharmingen during antibody development only or reported in the literature.

## Preparation and Storage

The monoclonal antibody was purified from tissue culture supernatant or ascites by affinity chromatography.

Store undiluted at 4°C.

## Application Notes

## Application

Flow cytometry	Routinely Tested
Immunohistochemistry-frozen	Tested During Development
Immunoprecipitation	Tested During Development

## Recommended Assay Procedure:

**Caution:** Sodium azide is a reversible inhibitor of oxidative metabolism; therefore, antibody preparations containing this preservative agent must not be used in cell cultures nor injected into animals. Sodium azide may be removed by washing stained cells or plate-bound antibody or dialyzing soluble antibody in sodium azide-free buffer. Since endotoxin may also affect the results of functional studies, we recommend the NA/LE™ (No Azide/Low Endotoxin) antibody format for in vitro and in vivo use.

## Suggested Companion Products

Catalog Number	Name	Size	Clone
554001	FITC Goat Anti-Mouse Ig	0.5 mg	Polyclonal

## Product Notices

1. Since applications vary, each investigator should titrate the reagent to obtain optimal results.
2. Please refer to [www.bdbiosciences.com/pharmingen/protocols](http://www.bdbiosciences.com/pharmingen/protocols) for technical protocols.
3. Caution: Sodium azide yields highly toxic hydrazoic acid under acidic conditions. Dilute azide compounds in running water before discarding to avoid accumulation of potentially explosive deposits in plumbing.

## References

Bevilacqua MP. Endothelial-leukocyte adhesion molecules. *Annu Rev Immunol.* 1993; 11:767-804. (Biology)

## BD Biosciences

[bdbiosciences.com](http://bdbiosciences.com)

United States	Canada	Europe	Japan	Asia Pacific	Latin America/Caribbean
877.232.8995	888.259.0187	32.53.720.550	0120.8555.90	65.6861.0633	55.11.5185.9995

For country-specific contact information, visit [bdbiosciences.com/how\\_to\\_order/](http://bdbiosciences.com/how_to_order/)

Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton Dickinson and Company is strictly prohibited.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.

BD, BD Logo and all other trademarks are the property of Becton, Dickinson and Company. ©2008 BD

