Technical Data Sheet

PE Rat Anti-Mouse CD210

Product	Information
---------	-------------

Material Number:	559914
Alternate Name:	IL-10 Receptor
Size:	0.2 mg
Concentration:	0.2 mg/ml
Clone:	1B1.3a
Immunogen:	Purified recombinant ligand-binding domain of mIL-10R
Isotype:	Rat IgG1, ĸ
Reactivity:	QC Testing: Mouse
Storage Buffer:	Aqueous buffered solution containing $\leq 0.09\%$ sodium azide.

Description

The 1B1.3a antibody reacts with the extracellular region of mouse CD210 which is also known as the mouse IL-10 receptor (mIL-10R); it does not recognize the human IL-10R. The IL-10R is expressed by a variety of mouse cell types and cell lines including thymocytes, T cells, B cells, and monocytes. 1B1.3a is a neutralizing antibody and reportedly blocks the binding of human IL-10, which cross-reacts with the mouse IL-10R. The immunogen used for the generation of the 1B1.3a hybridoma was purified recombinant ligand-binding domain of mIL-10R.



Mouse B220+ splenocytes were stained with R-PE-conjugated 1B1.3a (0.25 μg, Cat. No. 559914) (left panel). In addition, cells in buffer containing sodium azide were pre-incubated on ice with recombinant mouse IL-10 (0.25 μg, Cat. No. 550070) prior to staining with 1B1.3a to determine the blocking effect of bound IL-10 protein on staining with 1B1.3a (right panel). Staining with the R-PE-conjugated 1B1.3a antibody (filled histograms) is compared to the staining obtained using PE-conjugated R3-34 isotype control (0.25 μg, Cat. No. 554685) (open histograms). Histograms in both panels are gated on the lymphocyte population as defined by light scattering characteristics and were also gated on the cell subset stained with FITC-conjugated rat-anti-mouse B220 (0.06 μg, Cat. No. 553088).

Preparation and Storage

The monoclonal antibody was purified from tissue culture supernatant or ascites by affinity chromatography. The antibody was conjugated with R-PE under optimum conditions, and unconjugated antibody and free PE were removed. Store undiluted at 4° C and protected from prolonged exposure to light. Do not freeze.

BD Biosciences

www.bdbioscie	ences.com							
United States 877.232.8995 For country-spe	Canada 888.259.0187	Europe 32.53.720.550 ormation visit v	Japan 0120.8555.90 www.bdbioscience	Asia Pacific 65.6861.0633 es.com/how_to	Latin America/Caribbean 55.11.5185.9995 order/			
For country-specific contact information, visit www.bdbiosciences.com/how_to_order// Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product ther than the permitted use without the express written authorization of Becton Dickinson and Company is strictly prohibited. For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale. BD, BD Logo and all other trademarks are the property of Becton, Dickinson and Company. ©2007 BD								



Application Notes

Application

Flow cytometry Routinely Tested

Recommended Assay Procedure:

Immunofluorescent staining and flow cytometric analysis: The R-PE-conjugated 1B1.3a antibody (Cat. No. 559914) can be used for immmunofluorescent staining ($\leq 1 \mu g$ antibody/10e6 cells) and flow cytometric analysis of the levels of membrane IL-10R expressed by mouse cell lines or mouse leukocytes. An appropriate R-PE-conjugated immunoglobulin isotype-matched control is R3-34 (Cat. No. 554685).

Note: Since 1B1.3a is a neutralizing antibody, it competes with IL-10 for binding to its receptor. Therefore, the use of the 1B1.3a antibody for immunofluorescent staining and flow cytometric analysis in systems where the natural ligand of the receptor is present may give an underestimation of IL-10R expression. Based on our testing results, the presence of recombinant mouse IL-10 protein at levels above 250 ng/ml is sufficient to partially inhibit the binding of the 1B1.3a antibody (at 0.12 µg/10e6 cells).

In vitro neutralization: The NA/LE format of this antibody (Cat. No. 550012) is useful for bioassay.

Suggested Companion Products

Catalog Number	Name	Size	Clone	
554685	PE Rat IgG1, κ Isotype Control	0.1 mg	R3-34	
550070	Recombinant Mouse IL-10	10 µg	(none)	
553088	FITC Rat Anti-Mouse CD45R/B220	0.5 mg	RA3-6B2	

Product Notices

- 1. Since applications vary, each investigator should titrate the reagent to obtain optimal results.
- 2. Please refer to www.bdbiosciences.com/pharmingen/protocols for technical protocols.
- For fluorochrome spectra and suitable instrument settings, please refer to our Fluorochrome Web Page at www.bdbiosciences.com/pharmingen/colors.
- 4. Caution: Sodium azide yields highly toxic hydrazoic acid under acidic conditions. Dilute azide compounds in running water before discarding to avoid accumulation of potentially explosive deposits in plumbing.

References

O'Farrell AM, Liu Y, Moore KW, Mui AL. IL-10 inhibits macrophage activation and proliferation by distinct signaling mechanisms: evidence for Stat3-dependent and -independent pathways. *EMBO J.* 1998; 17(4):1006-1018.(Immunogen: Neutralization)