

Technical Data Sheet


Purified Rat Anti-Mouse CD195

Product Information

Material Number:	559921
Alternate Name:	CCR5
Size:	0.5 mg
Concentration:	0.5 mg/ml
Clone:	C34-3448
Immunogen:	Mouse CCR5 aa. 9-30
Isotype:	Rat IgG2c, κ
Reactivity:	QC Testing: Mouse
Storage Buffer:	Aqueous buffered solution containing $\leq 0.09\%$ sodium azide.

Description

The C34-3448 antibody reacts with mouse chemokine receptor CD195 (also known as CCR5), a member of the β -chemokine receptor family. CD195 regulates lymphocyte chemotaxis activation and transendothelial migration during inflammation. It signals a response to at least three chemokines: RANTES and macrophage inflammatory protein-1 (MIP-1) α and β . In the mouse system, the gene for CD195 has been mapped to chromosome 9. CD195 mRNA is expressed in heart, spleen and liver tissues, and by macrophages and T-lymphocytes. The immunogen used to generate the C34-3448 hybridoma was a KLH-conjugated peptide consisting of amino acids 9-30 of mouse CD195 (CCR5).



Expression of mouse CD195 by stimulated mouse splenocytes. BALB/c splenocytes were stimulated for 24 hours with ConA (10 μ g/ml, final concentration; Sigma). The splenocytes were cultured for additional 5 days in the presence of mouse IL-2 (100 U/ml, final concentration; Cat. No. 550069). The splenocytes were stained with purified rat anti-mouse CD195 antibody (C34-3448), followed by biotinylated mouse anti-rat IgG2c (A92-1, Cat. No. 553909) and Streptavidin-phycoerythrin (Cat. No. 554061) following BD Pharmingen's staining protocol (see Figure, Left panel). The data reflects gating on lymphocytes, based on forward and side scattered light signals. The level of nonspecific staining was assessed using the purified rat IgG2c isotype control (purified A23-1; Cat. No. 553982; Right panel). The quadrant markers for the bivariate dot plots were set based on the autofluorescence control and isotype control.

Preparation and Storage

The monoclonal antibody was purified from tissue culture supernatant or ascites by affinity chromatography.

Store undiluted at 4°C.

Application Notes

Application

Flow cytometry

Routinely Tested

Recommended Assay Procedure:

BD Biosciences

bdbiosciences.com

United States	Canada	Europe	Japan	Asia Pacific	Latin America/Caribbean
877.232.8995	888.259.0187	32.53.720.550	0120.8555.90	65.6861.0633	55.11.5185.9995

For country-specific contact information, visit bdbiosciences.com/how_to_order/

Conditions: The information disclosed herein is not to be construed as a recommendation to use the above product in violation of any patents. BD Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Purchase does not include or carry any right to resell or transfer this product either as a stand-alone product or as a component of another product. Any use of this product other than the permitted use without the express written authorization of Becton Dickinson and Company is strictly prohibited.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Not for resale.

BD, BD Logo and all other trademarks are the property of Becton, Dickinson and Company. ©2008 BD



Immunofluorescent Staining and Flow Cytometry Analysis: The purified C34-3448 antibody can be used for multicolor immunofluorescent staining and flow cytometric analysis to identify and enumerate CD195-expressing cells within mixed cell populations. For optimal immunofluorescent staining with flow cytometric analysis, the antibody should be titrated (≤ 0.25 μg mAb/million cells, see Figure above). For specific methodology, please visit the protocols section or chapter on intracellular staining in the Immune Function Handbook, both of which are posted on our web site, www.bdbiosciences.com.

Suggested Companion Products

Catalog Number	Name	Size	Clone
550069	Recombinant mouse IL-2	20 μg	(none)
553982	Purified Rat IgG2c, κ Isotype Control	0.5 mg	A23-1
553909	Biotin Mouse Anti-Rat IgG2c	0.5 mg	A92-1
554061	PE Streptavidin	0.5 mg	(none)

Product Notices

1. Since applications vary, each investigator should titrate the reagent to obtain optimal results.
2. Please refer to www.bdbiosciences.com/pharming/en/protocols for technical protocols.
3. Caution: Sodium azide yields highly toxic hydrazoic acid under acidic conditions. Dilute azide compounds in running water before discarding to avoid accumulation of potentially explosive deposits in plumbing.

References

Boring L, Gosling J, Monteclaro FS, Lulis AJ, Tsou CL, Charo IF. Molecular cloning and functional expression of murine JE (monocyte chemoattractant protein 1) and murine macrophage inflammatory protein 1alpha receptors: evidence for two closely linked C-C chemokine receptors on chromosome 9. *J Biol Chem.* 1996; 271(13):7551-7558. (Biology)

Meyer A, Coyle AJ, Proudfoot AE, Wells TN, Power CA. Cloning and characterization of a novel murine macrophage inflammatory protein-1 alpha receptor. *J Biol Chem.* 1996; 271(24):14445-14451. (Biology)

Napolitano M, Seamon KB, Leonard WJ. Identification of cell surface receptors for the Act-2 cytokine. *J Exp Med.* 1990; 172(1):285-289. (Biology)